

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
18 October 2001 (18.10.2001)

PCT

(10) International Publication Number
WO 01/77676 A1

(51) International Patent Classification⁷: G01N 33/53,
33/567, C07K 14/47, 16/18, 14/705, 14/72, A61K 48/00,
C12N 5/10, 15/12, 15/62, 15/63, 15/64

(21) International Application Number: PCT/US01/10739

(22) International Filing Date: 4 April 2001 (04.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/195,532 7 April 2000 (07.04.2000) US
60/247,014 13 November 2000 (13.11.2000) US

(71) Applicant: SENOMYX, INC. [US/US]; Suite 160, 11099
N. Torrey Pines Road, La Jolla, CA 92037 (US).

(72) Inventor: ADLER, Jon, Elliot; Apartment 15, 1099
Turquoise Street, San Diego, CA 92109 (US).

(74) Agents: TESKIN, Robin, L. et al.; Pillsbury Winthrop
LLP, 1600 Tysons Boulevard, McLean, VA 22102 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: T2R TASTE RECEPTORS AND GENES ENCODING SAME

(57) Abstract: Newly identified mammalian taste-cell-specific G protein-Coupled Receptors and the genes encoding said receptors are described. Specifically, T2R taste G Protein-Coupled Receptors that are believed to be involved in bitter taste sensation, and the genes encoding the same, are described, along with methods for isolating such genes and for isolating and expressing such receptors. Methods for representing taste perception of a particular tastant in a mammal are also described, as are methods for generating a novel molecules or combinations of molecules that elicit a predetermined taste perception in a mammal, and methods for simulating one or more tastes.

WO 01/77676 A1

T2R TASTE RECEPTORS AND GENES ENCODING SAMECross Reference to Related Applications

This application claims priority to U.S. Serial No. 60/195,532, filed April 7, 2000, and U.S. Serial No. 60/247,014, filed November 13, 2000, both of which are herein incorporated by reference in their entireties.

Field of the Invention

The invention relates to newly identified mammalian chemosensory G Protein-Coupled Receptors, to a family of such receptors, and to the genes and cDNA encoding said receptors. More particularly, the invention relates to newly identified mammalian chemosensory G Protein-Coupled Receptors active in taste signaling, to a family of such receptors, to the genes and cDNA encoding said receptors, and to methods of using such receptors, genes, and cDNA in the analysis and discovery of taste modulators.

Description of the Related Art

The taste system provides sensory information about the chemical composition of the external world. Taste transduction is one of the most sophisticated forms of chemical-triggered sensation in animals, and is found throughout the animal kingdom, from simple metazoans to the most complex of vertebrates. Mammals are believed to have five basic taste modalities: sweet, bitter, sour, salty, and umami (the taste of monosodium glutamate).

Each taste modality is believed to be mediated by distinct transduction pathways. These pathways are believed to be mediated by receptors, *e.g.*, metabotropic or ionotropic receptors, expressed in subsets of taste receptor cells. For instance, some tastes are believed to be mediated by G Protein-Coupled Receptors, while other tastes are believed to be mediated by channel proteins (*see, e.g.*, Kawamura *et al.*, *Introduction to Umami: A Basic Taste* (1987); Kinnamon *et al.*, *Ann. Rev. Physiol.*, 54:715-31 (1992); Lindemann, *Physiol. Rev.*, 76:718-66 (1996); Stewart *et al.*, *Am. J. Physiol.*, 272:1-26(1997)).

In mammals, taste receptor cells are assembled into taste buds that are distributed into different papillae in the tongue epithelium. Circumvallate papillae, found at the very back of the tongue, contain hundreds to thousands of taste buds. By contrast, foliate papillae, localized to the posterior lateral edge of the tongue, contain
5 dozens to hundreds of taste buds. Further, fungiform papillae, located at the front of the tongue, contain only a small number of taste buds.

Each taste bud, depending on the species, contains 50-150 cells, including precursor cells, support cells, and taste receptor cells. *See, e.g., Lindemann, Physiol. Rev., 76:718-66 (1996).* Receptor cells are innervated at their base by afferent nerve
10 endings that transmit information to the taste centers of the cortex through synapses in the brain stem and thalamus. Elucidating the mechanisms of taste cell signaling and information processing is important to understanding the function, regulation, and perception of the sense of taste.

Numerous physiological studies in animals have shown that taste receptor cells
15 may selectively respond to different chemical stimuli (*see, e.g., Akabas et al., Science, 242:1047-50 (1988); Gilbertson et al., J. Gen. Physiol., 100:803-24 (1992); Bernhardt et al., J. Physiol., 490:325-36 (1996); Cummings et al., J. Neurophysiol., 75:1256-63 (1996)*). More particularly, cells that express taste receptors, when exposed to certain chemical stimuli, elicit taste sensation by depolarizing to generate an action potential.
20 The action potential is believed to trigger the release of neurotransmitters at gustatory afferent neuron synapses, thereby initiating signaling along neuronal pathways that mediate taste perception (*see, e.g., Roper, Ann. Rev. Neurosci., 12:329-53 (1989)*). Nonetheless, at present, the means by which taste sensations are elicited remains poorly understood (*see, e.g., Margolskee, BioEssays, 15:645-50 (1993); Avenet et al.,*
25 *J. Membrane Biol., 112:1-8 (1989)*).

As described above, taste receptors specifically recognize molecules that elicit specific taste sensation. These molecules are also referred to herein as "tastants." Many taste receptors belong to the 7-transmembrane receptor superfamily (Hoon *et al., Cell* 96:451 (1999); Adler *et al., Cell*, 100:693 (2000)), which are also known as
30 G Protein-Coupled Receptors (GPCRs). G Protein-Coupled Receptors control many physiological functions, such as endocrine function, exocrine function, heart rate, lipolysis, and carbohydrate metabolism. The biochemical analysis and molecular

cloning of a number of such receptors has revealed many basic principles regarding the function of these receptors.

For example, U. S. Patent No. 5,691,188 describes how upon a ligand binding to a GPCR, the receptor presumably undergoes a conformational change leading to activation of the G Protein. G Proteins are comprised of three subunits: a guanyl nucleotide binding α subunit, a β subunit, and a γ subunit. G Proteins cycle between two forms, depending on whether GDP or GTP is bound to the α subunit. When GDP is bound, the G Protein exists as a heterotrimer: the $G\alpha\beta\gamma$ complex. When GTP is bound, the α subunit dissociates from the heterotrimer, leaving a $G\beta\gamma$ complex. When a $G\alpha\beta\gamma$ complex operatively associates with an activated G Protein-Coupled Receptor in a cell membrane, the rate of exchange of GTP for bound GDP is increased and the rate of dissociation of the bound $G\alpha$ subunit from the $G\alpha\beta\gamma$ complex increases. The free $G\alpha$ subunit and $G\beta\gamma$ complex are thus capable of transmitting a signal to downstream elements of a variety of signal transduction pathways. These events form the basis for a multiplicity of different cell signaling phenomena, including for example the signaling phenomena that are identified as neurological sensory perceptions such as taste and/or smell.

Complete or partial sequences of numerous human and other eukaryotic chemosensory receptors are currently known (*see, e.g.,* Pilpel, Y. *et al.*, *Protein Science*, 8:969-77 (1999); Mombaerts, P., *Annu. Rev. Neurosci.*, 22:487-50 (1999); EP0867508A2; US 5874243; WO 92/17585; WO 95/18140; WO 97/17444; WO 99/67282). Although much is known about the psychophysics and physiology of taste cell function, very little is known about the molecules and pathways that mediate its sensory signaling response. The identification and isolation of novel taste receptors and taste signaling molecules could allow for new methods of chemical and genetic modulation of taste transduction pathways. For example, the availability of receptor and channel molecules could permit screening for high affinity agonists, antagonists, inverse agonists, and modulators of taste activity. Such taste modulating compounds could be useful in the pharmaceutical and food industries to improve the taste of a variety of consumer products, or to block undesirable tastes, *e.g.* bitter tastes, in certain products.

Summary of the Invention

In part, the present invention addresses the need for better understanding of the interactions between chemosensory receptors and chemical stimuli. Thus, the present invention provides, among other things, novel taste receptors, and methods for
5 utilizing such receptors, and the genes and cDNAs encoding such receptors, to identify molecules that can be used to modulate taste transduction.

More particularly, the invention relates to a recently discovered family of G Protein-Coupled Receptors, and to the genes and cDNAs encoding said receptors. The receptors are thought to be primarily involved in bitter taste transduction, but can
10 be involved in transducing signals from other taste modalities as well.

The invention provides methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the receptors and genes encoding said receptors disclosed herein.

15 Toward that end, it is an object of the invention to provide a new family of mammalian G Protein-Coupled Receptors, herein referred to as T2Rs, active in taste perception. It is another object of the invention to provide fragments and variants of such T2Rs that retain tastant-binding activity.

It is yet another object of the invention to provide nucleic acid sequences or
20 molecules that encode such T2Rs, fragments, or variants thereof.

It is still another object of the invention to provide expression vectors which include nucleic acid sequences that encode such T2Rs, or fragments or variants thereof, which are operably linked to at least one regulatory sequence such as a promoter, enhancer, or other sequence involved in positive or negative gene
25 transcription and/or translation.

It is still another object of the invention to provide human or non-human cells that functionally express at least one of such T2Rs, or fragments or variants thereof.

It is still another object of the invention to provide T2R fusion proteins or polypeptides that include at least a fragment of at least one of such T2Rs.

30 It is another object of the invention to provide an isolated nucleic acid molecule encoding a T2R polypeptide comprising a nucleic acid sequence that is at least 50%, preferably 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a

nucleic acid sequence selected from the group consisting of: SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, and conservatively modified variants thereof.

It is a further object of the invention to provide an isolated nucleic acid molecule comprising a nucleic acid sequence that encodes a polypeptide having an amino acid sequence at least 35 to 50%, and preferably 60%, 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to an amino acid sequence selected from the group consisting of: SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, and conservatively modified variants thereof, wherein the fragment is at least 20, preferably 40, 60, 80, 100, 150, 200, or 250 amino acids in length. Optionally, the fragment can be an antigenic fragment that binds to an anti-T2R antibody.

It is still a further object of the invention to provide an isolated polypeptide comprising a variant of said fragment, wherein there is a variation in at most 10, preferably 5, 4, 3, 2, or 1 amino acid residues.

It is still another object of the invention to provide agonists or antagonists of such T2Rs, or fragments or variants thereof.

It is yet another object of the invention to provide methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the T2Rs, or fragments or variants thereof, and genes encoding such T2Rs, or fragments or variants thereof, disclosed herein.

It is yet another object of the invention to provide novel molecules or combinations of molecules that elicit a predetermined taste perception in a mammal. Such molecules or compositions can be generated by determining a value of taste perception in a mammal for a known molecule or combinations of molecules; determining a value of taste perception in a mammal for one or more unknown molecules or combinations of molecules; comparing the value of taste perception in a mammal for one or more unknown compositions to the value of taste perception in a mammal for one or more known compositions; selecting a molecule or combination of molecules that elicits a predetermined taste perception in a mammal; and combining two or more unknown molecules or combinations of molecules to form a molecule or combination of molecules that elicits a predetermined taste perception in a mammal.

The combining step yields a single molecule or a combination of molecules that elicits a predetermined taste perception in a mammal.

It is still a further object of the invention to provide a method of screening one or more compounds for the presence of a taste detectable by a mammal, comprising: a
5 step of contacting said one or more compounds with at least one of the disclosed T2Rs, fragments or variants thereof, preferably wherein the mammal is a human.

It is another object of the invention to provided a method for simulating a taste, comprising the steps of: for each of a plurality of T2Rs, or fragments of variants thereof disclosed herein, preferably human T2Rs, ascertaining the extent to which the
10 T2R interacts with the tastant; and combining a plurality of compounds, each having a previously ascertained interaction with one or more of the T2Rs, in amounts that together provide a receptor-stimulation profile that mimics the profile for the taste. Interaction of a tastant with a T2R can be determined using any of the binding or reporter assays described herein. The plurality of compounds may then be combined
15 to form a mixture. If desired, one or more of the plurality of the compounds can be combined covalently. The combined compounds substantially stimulate at least 50%, 60%, 70%, 75%, 80% or 90% or all of the receptors that are substantially stimulated by the tastant.

In yet another aspect of the invention, a method is provided wherein a plurality
20 of standard compounds are tested against a plurality of T2Rs, or fragments or variants thereof, to ascertain the extent to which the T2Rs each interact with each standard compound, thereby generating a receptor stimulation profile for each standard compound. These receptor stimulation profiles may then be stored in a relational database on a data storage medium. The method may further comprise providing a
25 desired receptor-stimulation profile for a taste; comparing the desired receptor stimulation profile to the relational database; and ascertaining one or more combinations of standard compounds that most closely match the desired receptor-stimulation profile. The method may further comprise combining standard compounds in one or more of the ascertained combinations to simulate the taste.

30 It is a further object of the invention to provide a method for representing taste perception of a particular substance in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n T2Rs of said

vertebrate, where n is greater than or equal to 4, n is greater than or equal to 12; n is greater than or equal to 24, or n is greater than or equal to 40; and generating from said values a quantitative representation of taste perception. The T2Rs may be a taste receptor disclosed herein, or fragments or variants thereof, the representation may constitute a point or a volume in n -dimensional space, may constitute a graph or a spectrum, or may constitute a matrix of quantitative representations. Also, the providing step may comprise contacting a plurality of recombinantly produced T2Rs, or fragments or variants thereof, with a test composition and quantitatively measuring the interaction of said composition with said receptors.

10 It is a related object of the invention to provide a method for predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n T2Rs of said vertebrate, where n is greater than or equal to 4 n is greater than or equal to 12; n is greater than or equal to 24, or n is greater than or equal to 40; for one or more molecules or combinations of molecules yielding known taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal, providing values X_1 to X_n representative of the quantitative stimulation of each of n T2Rs of said vertebrate, where n is greater than or equal to 4, n is greater than or equal to 12; n is greater than or equal to 24, or n is greater than or equal to 40; for one or more molecules or combinations of molecules yielding unknown taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, and predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal by comparing the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal to the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal. The T2Rs

used in this method may include a taste receptor, or fragment or variant thereof, disclosed herein.

Detailed Description of the Invention

5 The invention thus provides isolated nucleic acid molecules encoding taste-cell-specific G Protein-Coupled Receptors ("GPCRs"), and the polypeptides they encode. These nucleic acid molecules and the polypeptides that they encode are members of the T2R family of taste-cell-specific GPCRs. More particularly, the recent identification of the T2R gene family, which encodes candidate bitter taste
10 receptors, database accession numbers AF227129-AF227149 and AF240765-AF240768, prompted the search for, and identification of related genes in public nucleotide sequence databases. The present invention relates to newly identified members of this family. Further information regarding the T2R family can be found in Adler *et al.*, *Cell*, 100:693-702 (2000) and Chandrashekar *et al.*, *Cell*, 100:703-11
15 (200), both of which are herein incorporated by reference in their entireties.

 Nucleic acids encoding the T2R proteins and polypeptides of the invention can be isolated from a variety of sources, genetically engineered, amplified, synthesized, and/or expressed recombinantly according to the methods disclosed in WO 0035374, which is herein incorporated by reference in its entirety.

20 More particularly, the invention provides nucleic acids encoding a novel family of taste-cell-specific G Protein-Coupled Receptors. These nucleic acids and the receptors that they encode are referred to as members of the "T2R" family of taste-cell-specific G Protein-Coupled Receptors ("GPCRs"). These taste-cell-specific GPCRs are believed to be components of the taste transduction pathway, specifically,
25 the bitter taste transduction pathway, and are involved in the taste detection of substances such as the bitter substances, 6-n-propylthiouracil (PROP), sucrose octaacetate (soa), raffinose undecaacetate (rua), denatonium, copper glycinate, and quinine. However, the T2Rs may be involved in other taste modalities as well.

 Further, it is believed that T2R family members may act in combination with
30 other T2R family members, other taste-cell-specific GPCRs, or a combination thereof, to thereby effect chemosensory taste transduction. For instance, it is believed that T2R family members maybe co-expressed within the same taste receptor cell type, and

the co-expressed receptors may physically interact to form a heterodimeric taste receptor. Alternatively, the co-expressed receptors may both independently bind to the same type of ligand, and their combined binding may result in a specific perceived taste sensation.

5 The invention also provides methods of screening for modulators, *e.g.*, activators, inhibitors, stimulators, enhancers, agonists, and antagonists, of these novel taste-cell-specific GPCRs. Such modulators of taste transduction are useful for pharmacological and genetic modulation of taste signaling pathways. These methods of screening can be used to identify high affinity agonists and antagonists of taste cell
10 activity. These modulatory compounds can then be used in the food and pharmaceutical industries to customize taste, for example, to decrease or mask the bitter taste of foods or drugs.

Thus, the invention provides assays for taste modulation, where members of the T2R family act as direct or indirect reporter molecules of the effect of modulators
15 on taste transduction. GPCRs can be used in assays, *e.g.*, to measure changes in ligand binding, ion concentration, membrane potential, current flow, ion flux, transcription, signal transduction, receptor-ligand interactions, second messenger concentrations, *in vitro*, *in vivo*, and *ex vivo*. In one embodiment, members of the T2R family can be used as indirect reporters via attachment to a second reporter
20 molecule such as green fluorescent protein (*see, e.g.*, Mistili & Spector, *Nature Biotechnology*, 15:961-64 (1997)). In another embodiment, T2R family members are recombinantly expressed in cells, and modulation of taste transduction via GPCR activity can be assayed by measuring changes in Ca^{2+} levels and other intracellular messengers such as cAMP, cGMP, or IP3.

25 In a preferred embodiment, a T2R polypeptide is expressed in an eukaryotic cell as a chimeric receptor with a heterologous sequence that facilitates plasma membrane trafficking or maturation and targeting through the secretory pathway. In a preferred embodiment, the heterologous sequence is a rhodopsin sequence, such as an N-terminal fragment of a rhodopsin. Such chimeric T2R receptors can be expressed
30 in any eukaryotic cell, such as HEK-293 cells. Preferably, the cells comprise a functional G Protein, *e.g.*, G α 15, that is capable of coupling the chimeric receptor to an intracellular signaling pathway or to a signaling protein such as phospholipase C.

Activation of such chimeric receptors in such cells can be detected using any standard method, such as by detecting changes in intracellular calcium by detecting FURA-2-dependent fluorescence in the cell. If host cells do not express an appropriate G Protein, they may be transfected with a gene encoding a promiscuous G Protein such as those described in US Serial No. 60/243,770, which is herein incorporated by reference in its entirety.

Methods of assaying for modulators of taste transduction can include *in vitro* ligand binding assays using T2R polypeptides, portions thereof, such as the extracellular domain, transmembrane region, or combinations thereof, or chimeric proteins comprising one or more domains of a T2R family member; oocyte, primary cell or tissue culture cell T2R gene expression, or expression of T2R fragments or fusion proteins, such as rhodopsin fusion proteins; phosphorylation and dephosphorylation of T2R family members; G Protein binding to GPCRs; arrestin binding; internalization; ligand binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cGMP, cAMP and IP₃; changes in intracellular Ca²⁺ levels; and neurotransmitter release.

Further, the invention provides methods of detecting T2R nucleic acid and protein expression, allowing investigation of taste transduction regulation and specific identification of taste receptor cells. T2R family members also provide useful nucleic acid probes for paternity and forensic investigations. T2R genes are also useful as a nucleic acid probes for identifying taste receptor cells, such as foliate, fungiform, circumvallate, geschmackstreifen, and epiglottis taste receptor cells, in particular bitter-taste receptive, gustducin expressing cells. Furthermore, the nucleic acids and the polypeptides they encode can be used as probes to dissect taste-induced behaviors.

T2R polypeptides can also be used to generate monoclonal and polyclonal antibodies useful for identifying taste receptor cells. Taste receptor cells can be identified using techniques such as reverse transcription and amplification of mRNA, isolation of total RNA or poly A⁺ RNA, northern blotting, dot blotting, *in situ* hybridization, RNase protection, S1 digestion, probing DNA microchip arrays, western blots, and the like.

The T2R genes and the polypeptides they encode comprise a family of related taste-cell-specific G Protein-Coupled Receptors. Within the genome, these genes are present either alone or within one of several gene clusters. One gene cluster, located at human genomic region 12p13, comprises at least 24 genes, and a second cluster, located at 7q33, comprises at least 7 genes. In total, more than 60 distinct T2R family members have been identified from difference organisms, including several putative pseudogenes. It is estimated that the human genome may include approximately 40 distinct T2R genes, encoding about 30 functional human receptors.

Some of the T2R genes have been associated with previously mapped loci implicated in the control of bitter taste. For example, the human T2R1 is located at human interval 5p15, precisely where a locus that influences the ability to taste the substance PROP has previously been mapped (*see Reed et al., Am. J. Hum. Genet.*, 64:1478-80 (1999)). In addition, the human gene cluster found at genomic region 12p13 corresponds to a region of mouse chromosome 6 that has been shown to contain numerous bitter-tasting genes, that are believed to influence taste perception or sensation including *e.g.*, sucrose octaacetate, raffinose, undecaacetate cycloheximide, and quinine (*see, e.g., Lush et al., Genet. Res.*, 6:167-74 (1995)). These associations suggest that the T2R genes are involved in the taste detection of various substances, in particular bitter substances. In addition, mouse T2R5 is specifically receptive to cycloheximide, and mutations in the mT2R5 gene have been hypothesized to produce a cycloheximide-non-tasting phenotype. Similarly, human T2R4 and mouse T2R8 are specifically responsive to both denatonium and PROP.

Functionally, the T2R genes comprise a family of related seven transmembrane G Protein-Coupled Receptors which are believed to be involved in taste transduction and which may interact with a G Protein to mediate taste signal transduction (*see, e.g., Fong, Cell Signal*, 8:217 (1996); Baldwin, *Curr. Opin. Cell Biol.*, 6:180 (1994)). In particular, T2Rs are believed to interact in a ligand-specific manner with the G Protein gustducin.

Structurally, the nucleotide sequences of T2R family members encode a family of related polypeptides comprising an extracellular domain, seven transmembrane domains, and a cytoplasmic domain. Related T2R family genes from other species typically will share about 20-30% nucleotide sequence identity over a region of at

least about 50 nucleotides in length, optionally 100, 200, 500, or more nucleotides in length to SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or conservatively modified variants thereof; or encode polypeptides sharing at least about 30-40% amino acid sequence identity over an amino acid region at least about 25 amino acids in length, optionally 50 to 100 amino acids in length to SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or conservatively modified variants thereof. It has been shown that T2R genes are selectively expressed in subsets of taste receptor cells of the tongue, palate epithelium, foliate, geschmackstreifen, and epiglottis. In contrast, studies have shown that T2Rs are less often expressed in fungiform papillae. Further, it has been shown that T2Rs are selectively expressed in gustducin-positive cells.

Several consensus amino acid sequences or domains have also been identified that are characteristic of T2R family members. Particularly, it has been found that T2R family members typically comprise a sequence having at least about 50%, optionally 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or higher, identity to T2R Transmembrane Region I, T2R Transmembrane Region II, T2R Transmembrane Region III, T2R Transmembrane Region IV, T2R Transmembrane Region V, and T2R Transmembrane Region VII. These conserved domains thus can be used to identify members of the T2R family, by identity, specific hybridization or amplification, or specific binding by antibodies raised against a domain. Such T2R transmembrane regions have the following amino acid sequences:

T2R Family Consensus Sequence 1:

E(F/A)(I/V/L)(V/L)G(I/V)(L/V)GN(G/T)FI(V/A)LVNC(I/M)DW (SEQ ID NO: 25)

T2R Family Consensus Sequence 2:

(D/G)(F/L)(I/L)L(T/I)(G/A/S)LAISRI(C/G/F)L (SEQ ID NO: 26)

T2R Family Consensus Sequence 3:

NH(L/F)(S/T/N)(L/I/V)W(F/L)(A/T)T(C/S/N)L(S/N/G)(I/V) (SEQ ID NO: 27)

T2R Family Consensus Sequence 4:

FY(F/C)LKIA(N/S)FS(H/N)(P/S)(L/I/V)FL(W/Y)LK (SEQ ID NO: 28)

T2R Family Consensus Sequence 5:

LLI(I/F/V)SLW(K/R)H(S/T)(K/R)(Q/K)(M/I)(Q/K) (SEQ ID NO: 29)

T2R Family Consensus Sequence 6:

HS(F/L)(I/V)LI(L/M)(G/S/T)N(P/S/N)KL(K/R)(Q/R) (SEQ ID NO: 30)

Specific regions of human T2R nucleotide and amino acid sequences may be used to identify polymorphic variants or alleles or homologs comprised in humans or other species. This identification can be effected *in vitro*, *e.g.*, under stringent hybridization conditions or PCR (*e.g.*, using primers encoding the T2R consensus sequences identified above) and sequencing, or by using the sequence information in a computer system for comparison with other nucleotide sequences. Typically, identification of polymorphic variants or alleles of T2R family members can be made by comparing an amino acid sequence of about 25 amino acids or more, *e.g.*, 50-100 amino acids. Amino acid identity of approximately 30-40%, optionally 50-60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95-99% or above suggests that a protein is a polymorphic variant, or allele, or a homolog or ortholog of a T2R family member. Sequence identity can be calculated by the methods disclosed *infra*. It can be determined almost definitely that a positive T2R gene falls within the T2R family based on the additional presence of at least one conserved sequence that is identical or at least 75% identical to the previously identified T2R family consensus sequences. Sequence comparison can be performed using any of the sequence comparison algorithms discussed below. Antibodies that bind specifically to T2R polypeptides or a conserved region thereof can also be used to identify alleles, interspecies homologs, and polymorphic variants of T2R proteins.

Polymorphic variants, or alleles, and related T2R homologs or orthologs can be confirmed by examining taste-cell-specific expression of the putative T2R polypeptide. Typically, T2R polypeptides having an amino acid sequence disclosed herein can be used as a positive control in comparison to the putative T2R protein to demonstrate the identification of an allele or homolog of the T2R family member. Such homologs or alleles will similarly possess the seven transmembrane structure of a G Protein-Coupled Receptor.

Nucleotide and amino acid sequence information for T2R family members may also be used to construct models of taste-cell-specific polypeptides in a computer system. These models are subsequently used to identify compounds that can activate or inhibit T2R receptor proteins. Such compounds that modulate the activity of T2R family members can be used to investigate the role of T2R genes and polypeptides in taste transduction.

The isolation of T2R family members provides a means for assaying for inhibitors and activators of taste transduction. Biologically active T2R proteins are useful for testing inhibitors and activators of T2R as taste transducers, especially bitter taste transducers, using *in vivo* (animal based assays) and *in vitro* assays that measure, 5 *e.g.*, ligand binding; phosphorylation and dephosphorylation; binding to G Proteins; G Protein activation; regulatory molecule binding; voltage, membrane potential and conductance changes; ion flux; intracellular second messengers such as cGMP, cAMP, IP3; and intracellular calcium levels. Such activators and inhibitors identified using T2R family members can be used to further study taste transduction and to 10 identify specific taste agonists and antagonists. Such activators and inhibitors are useful as pharmaceutical and food agents for customizing taste, for example to decrease the bitter taste of foods or pharmaceuticals.

The present invention also provides assays, preferably high throughput assays, to identify molecules that interact with and/or modulate a T2R polypeptide. In 15 numerous assays, a particular portion of a T2R family member will be used, *e.g.*, an extracellular, transmembrane, or intracellular or region. In numerous embodiments, an extracellular domain, transmembrane region, or combination thereof is bound to a solid substrate, and used, *e.g.*, to isolate ligands, agonists, inverse agonists, antagonists, or any other molecules that can bind to and/or modulate the activity of an 20 extracellular or transmembrane region of a T2R polypeptide.

In certain embodiments, a region of a T2R polypeptide, *e.g.*, an extracellular, transmembrane, or intracellular region, is fused to a heterologous polypeptide, thereby forming a chimeric polypeptide, *e.g.*, a chimeric polypeptide with GPCR activity. Such chimeric polypeptides are useful, *e.g.*, in assays to identify ligands, agonists, 25 inverse agonists, antagonists, or other modulators of a T2R polypeptide. In addition, such chimeric polypeptides are useful to create novel taste receptors with novel ligand binding specificity, modes of regulation, signal transduction pathways, or other such properties, or to create novel taste receptors with novel combinations of ligand binding specificity, modes of regulation, signal transduction pathways, etc. Also T2R 30 nucleic acids and expression of T2R polypeptides can be used to create topological maps of the tongue that potentially can be used to study the relation of tongue taste receptor cells to taste sensory neurons in the brain. In particular, methods of detecting

T2Rs can potentially be used to identify taste cells that are sensitive to bitter tasting substances. Chromosome localization of the genes encoding human T2R genes can also potentially be used to identify diseases, mutations, and traits caused by, or associated with T2R family members.

5 Generally, the invention provides isolated nucleic acid molecules of the T2R gene family and the taste receptors they encode. The present invention also includes not only nucleic acid and polypeptide sequences having the specified amino acid sequences, but also fragments, particularly fragments of, for example, 40, 60, 80, 100, 150, 200, or 250 nucleotides, or more, as well as protein fragments of, for example,
10 10, 20, 30, 50, 70, 100, or 150 amino acids, or more.

 Various conservative mutations and substitutions are envisioned to be within the scope of the invention. For instance, it would be within the level of skill in the art to perform amino acid substitutions using known protocols of recombinant gene technology including PCR, gene cloning, site-directed mutagenesis of cDNA,
15 transfection of host cells, and in-vitro transcription. The variants can then be screened for functional activity.

 More particularly, specific regions of the nucleic acid sequences disclosed herein, and the polypeptides they encode, may be used to identify polymorphic variants, interspecies homologs, and alleles of the sequences. This identification can
20 be made *in vitro*, e.g., under stringent hybridization conditions, PCR, and sequencing, or by using the sequence information in a computer system for comparison with other nucleic acid sequences. Different alleles of T2R genes within a single species population will also be useful in determining whether differences in allelic sequences correlate to differences in taste perception between members of the population.

25 The nucleic acid molecules of the present invention are generally intronless and encode putative T2R proteins generally having lengths of approximately 300 residues and are comprised of seven transmembrane regions, as predicted by hydrophobicity plotting analysis, suggesting that they belong to the G Protein-Coupled Receptor (7TM) superfamily. In addition to the overall structural similarity, each of
30 the T2Rs identified herein has a characteristic sequence signature of a T2R family member. In particular, all sequences contain very close matches to the T2R family consensus sequences identified above. Combination of all the above mentioned

structural features of the identified genes and encoded proteins strongly suggests that they represent novel members of the T2R receptor family.

It is also hypothesized that that T2R receptors and their genes can be used, alone or in combination with other types of taste receptors, in developing detection systems and assays for chemically identifying distinct types of molecules specifically
5 recognized by these receptors, as well as for diagnostic and research purposes.

The nucleic acid sequences of the invention and other nucleic acids used to practice this invention, whether RNA, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered,
10 amplified, and/or expressed recombinantly. Any recombinant expression system can be used, including, in addition to mammalian cells, *e.g.*, bacterial, yeast, insect, or plant systems.

A. Identification of T2Rs

The amino acid sequences of the T2R proteins and polypeptides of the invention can be identified by putative translation of the coding nucleic acid sequences. These various amino acid sequences and the coding nucleic acid sequences may be compared to one another or to other sequences according to a number of methods. In a particular embodiment, the pseudogenes disclosed herein
20 can be used to identify functional alleles or related genes in genomic databases known in the art.

For example, in sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm
25 program parameters are designated. Default program parameters can be used, as described below for the BLASTN and BLASTP programs, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on
30 the program parameters.

A "comparison window," as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of

from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.*, 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *PNAS*, 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds. 1995 supplement)).

A preferred example of an algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.*, *Nuc. Acids Res.*, 25:3389-402 (1977) and Altschul *et al.*, *J Mol. Biol.*, 215:403-10 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al.*, *Nuc. Acids Res.*, 25:3389-402 (1977) and Altschul *et al.*, *J Mol. Biol.*, 215:403-10 (1990)). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or

below, due to the accumulation of one or more negative scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (*see* Henikoff & Henikoff, *PNAS*, 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

Another example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a so-called "tree" or "dendrogram" showing the clustering relationships used to create the alignment (*see, e.g.*, Figure 2). PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J Mol. Evol.*, 35:351-60 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151-53 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, *e.g.*, version 7.0 Devereaux *et al.*, *Nuc. Acids Res.*, 12:387-95 (1984). Comparison of these protein sequences to all known proteins in the public sequence databases using BLASTP algorithm revealed their strong homology to the members of the T2R family, each of

the T2R family sequences having at least 50%, and preferably at least 55%, at least 60%, at least 65%, and most preferably at least 70%, amino acid identity to at least one known member of the family.

5 **B. Definitions**

As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

“Taste cells” include neuroepithelial cells that are organized into groups to form taste buds of the tongue, *e.g.*, foliate, fungiform, and circumvallate cells (*see, e.g.*, Roper *et al.*, *Ann. Rev. Neurosci.* 12:329-353 (1989)). Taste cells are also found
10 in the palate, and other tissues such as the esophagus and the stomach.

“T2R” refers to one or more members of a family of G Protein-Coupled Receptors that are selectively expressed in taste cells of the tongue and palate epithelium, such as foliate, geschmackstreifen, epiglottis, fungiform, and
15 circumvallate cells, as well as cells of the esophagus, and stomach (*see, e.g., Adler et al., Cell*, 100:693-702 (2000)). This family is also referred to as the “SF family” (*see, e.g., PCT/US00/24821*, which is herein incorporated by reference in its entirety). Such taste cells can be identified because they express specific molecules such as gustducin, a taste-cell-specific G Protein, or other taste specific molecules
20 (McLaughlin *et al., Nature*, 357:563-69 (1992)). Taste receptor cells can also be identified on the basis of morphology (*see, e.g., Roper, supra*). T2R family members have the ability to act as receptors for taste transduction. T2R family members are also referred to as the “GR” family, for gustatory receptor, or “SF” family.

“T2R” nucleic acids encode a family of GPCRs with seven transmembrane
25 regions that have “G Protein-Coupled Receptor activity,” *e.g.*, they may bind to G Proteins in response to extracellular stimuli and promote production of second messengers such as IP3, cAMP, cGMP, and Ca²⁺ via stimulation of enzymes such as phospholipase C and adenylate cyclase (for a description of the structure and function of GPCRs, *see, e.g., Fong, supra*, and Baldwin, *supra*). These nucleic acids encode
30 proteins that are expressed in taste cells, in particular gustducin-expressing taste cells that are responsive to bitter tastants. A single taste cell may contain many distinct T2R polypeptides.

The term "T2R" family therefore includes polymorphic variants, alleles, mutants, and homologs that: (1) have about 30-40% amino acid sequence identity, more specifically about 40, 50, 60, 70, 75, 80, 85, 90, 95, 96, 97, 98, or 99% amino acid sequence identity to SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, over
5 a window of about 25 amino acids, optionally 50-100 amino acids; (2) specifically bind to antibodies raised against an immunogen comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, and conservatively modified variants thereof; (3) specifically hybridize (with a size of at least about 100, optionally at least about 500-1000 nucleotides) under
10 stringent hybridization conditions to a sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, and conservatively modified variants thereof; (4) comprise a sequence at least about 40% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24; or (5) are amplified by primers that specifically hybridize under
15 stringent hybridization conditions to the same sequence as a degenerate primer sets encoding SEQ ID NOS: 25-30.

As previously discussed, while T2R genes exhibit substantial sequence divergence at both the protein and DNA level, all isolated to date have been found to contain comprise certain consensus sequences, in particular regions that are identical
20 or which possess or at least 70-75% sequence identity to the T2R consensus sequence identified previously (SEQ ID NOS: 25-30).

Topologically, certain chemosensory GPCRs have an "N-terminal domain;" "extracellular domains," a "transmembrane domain" comprising seven transmembrane regions, and corresponding cytoplasmic and extracellular loops,
25 "cytoplasmic regions," and a "C-terminal region" (*see, e.g., Hoon et al., Cell, 96:541-51 (1999); Buck & Axel, Cell, 65:175-87 (1991)*). These regions can be structurally identified using methods known to those of skill in the art, such as sequence analysis programs that identify hydrophobic and hydrophilic domains (*see, e.g., Stryer, Biochemistry, (3rd ed. 1988); see also any of a number of Internet based sequence*
30 *analysis programs, such as those found at dot.imgen.bcm.tmc.edu*). These regions are useful for making chimeric proteins and for in vitro assays of the invention, *e.g., ligand binding assays.*

“Extracellular domains” therefore refers to the domains of T2R polypeptides that protrude from the cellular membrane and are exposed to the extracellular face of the cell. Such regions would include the “N-terminal domain” that is exposed to the extracellular face of the cell, as well as the extracellular loops of the transmembrane domain that are exposed to the extracellular face of the cell, *i.e.*, the extracellular loops between transmembrane regions 2 and 3, transmembrane regions 4 and 5, and transmembrane regions 6 and 7. The “N-terminal domain” starts at the N-terminus and extends to a region close to the start of the transmembrane region. These extracellular regions are useful for *in vitro* ligand binding assays, both soluble and solid phase. In addition, transmembrane regions, described below, can also be involved in ligand binding, either in combination with the extracellular region or alone, and are therefore also useful for *in vitro* ligand binding assays.

“Transmembrane domain,” which comprises the seven transmembrane “regions,” refers to the domain of T2R polypeptides that lies within the plasma membrane, and may also include the corresponding cytoplasmic (intracellular) and extracellular loops, also referred to as transmembrane “regions.” The seven transmembrane regions and extracellular and cytoplasmic loops can be identified using standard methods, as described in Kyte & Doolittle, *J. Mol. Biol.*, 157:105-32 (1982)), or in Stryer, *supra*.

“Cytoplasmic domains” refers to the domains of T2R proteins that face the inside of the cell, *e.g.*, the “C-terminal domain” and the intracellular loops of the transmembrane domain, *e.g.*, the intracellular loops between transmembrane regions 1 and 2, transmembrane regions 3 and 4, and transmembrane regions 5 and 6. “C-terminal domain” refers to the region that spans from the end of the last transmembrane region to the C-terminus of the protein, and which is normally located within the cytoplasm.

The term “7- transmembrane receptor” means a polypeptide belonging to a superfamily of transmembrane proteins that have seven regions that span the plasma membrane seven times (thus, the seven regions are called “transmembrane” or “TM” domains TM I to TM VII). The families of olfactory and certain taste receptors each belong to this super-family. 7-transmembrane receptor polypeptides have similar and

characteristic primary, secondary and tertiary structures, as discussed in further detail below.

The term "ligand-binding region" refers to sequences derived from a chemosensory or taste receptor that substantially incorporates transmembrane domains II to VII (TM II to VII). The region may be capable of binding a ligand, and more particularly, a tastant.

The term "plasma membrane translocation domain" or simply "translocation domain" means a polypeptide domain that is functionally equivalent to an exemplary translocation domain (5'-MNGTEGPNFYVPFSNKTGVV; SEQ ID NO: 31). These peptide domains, when incorporated into the amino terminus of a polypeptide coding sequence, can with great efficiency "chaperone" or "translocate" the hybrid ("fusion") protein to the cell plasma membrane. This particular "translocation domain" was initially derived from the amino terminus of the human rhodopsin receptor polypeptide, a 7-transmembrane receptor. Another translocation domain has been derived from the bovine rhodopsin sequence and is also useful for facilitating translocation. Rhodopsin derived sequences are particularly efficient in translocating 7-transmembrane fusion proteins to the plasma membrane.

"Functional equivalency" means the domain's ability and efficiency in translocating newly translated proteins to the plasma membrane as efficiently as exemplary SEQ ID NO: 31 under similar conditions; relative efficiencies can be measured (in quantitative terms) and compared, as described herein. Domains falling within the scope of the invention can be determined by routine screening for their efficiency in translocating newly synthesized polypeptides to the plasma membrane in a cell (mammalian, *Xenopus*, and the like) with the same efficiency as the twenty amino acid long translocation domain SEQ ID NO: 31.

The phrase "functional effects" in the context of assays for testing compounds that modulate T2R family member mediated taste transduction includes the determination of any parameter that is indirectly or directly under the influence of the receptor, *e.g.*, functional, physical and chemical effects. It includes ligand binding, changes in ion flux, membrane potential, current flow, transcription, G Protein binding, GPCR phosphorylation or dephosphorylation, signal transduction, receptor-ligand interactions, second messenger concentrations (*e.g.*, cAMP, cGMP, IP3, or

intracellular Ca^{2+}), *in vitro*, *in vivo*, and *ex vivo* and also includes other physiologic effects such increases or decreases of neurotransmitter or hormone release.

By "determining the functional effect" is meant assays for a compound that increases or decreases a parameter that is indirectly or directly under the influence of a T2R family member, *e.g.*, functional, physical and chemical effects. Such functional effects can be measured by any means known to those skilled in the art, *e.g.*, changes in spectroscopic characteristics (*e.g.*, fluorescence, absorbance, refractive index), hydrodynamic (*e.g.*, shape), chromatographic, or solubility properties, patch clamping, voltage-sensitive dyes, whole cell currents, radioisotope efflux, inducible markers, oocyte T2R gene expression; tissue culture cell T2R expression; transcriptional activation of T2R genes; ligand binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cAMP, cGMP, and inositol triphosphate (IP3); changes in intracellular calcium levels; neurotransmitter release, and the like.

"Inhibitors," "activators," and "modulators" of T2R genes or proteins are used interchangeably to refer to inhibitory, activating, or modulating molecules identified using *in vitro* and *in vivo* assays for taste transduction, *e.g.*, ligands, agonists, antagonists, and their homologs and mimetics. Inhibitors are compounds that, *e.g.*, bind to, partially or totally block stimulation, decrease, prevent, delay activation, inactivate, desensitize, or down regulate taste transduction, *e.g.*, antagonists. Activators are compounds that, *e.g.*, bind to, stimulate, increase, open, activate, facilitate, enhance activation, sensitize, or up regulate taste transduction, *e.g.*, agonists. Modulators include compounds that, *e.g.*, alter the interaction of a receptor with: extracellular proteins that bind activators or inhibitor (*e.g.*, ebnerin and other members of the hydrophobic carrier family); G Proteins; kinases (*e.g.*, homologs of rhodopsin kinase and beta adrenergic receptor kinases that are involved in deactivation and desensitization of a receptor); and arrestins, which also deactivate and desensitize receptors. Modulators include genetically modified versions of T2R family members, *e.g.*, with altered activity, as well as naturally occurring and synthetic ligands, antagonists, agonists, small chemical molecules and the like.

Such assays for inhibitors and activators include, *e.g.*, expressing T2R family members in cells or cell membranes, applying putative modulator compounds in the

presence or absence of tastants, *e.g.*, bitter tastants, and then determining the functional effects on taste transduction, as described above. Samples or assays comprising T2R family members that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of modulation. Control samples (untreated with modulators) are assigned a relative T2R activity value of 100%. Inhibition of a T2R is achieved when the T2R activity value relative to the control is about 80%, optionally 50% or 25-0%. Activation of a T2R is achieved when the T2R activity value relative to the control is 110%, optionally 150%, optionally 200-500%, or 1000-3000% higher.

The terms "purified," "substantially purified," and "isolated" as used herein refer to the state of being free of other, dissimilar compounds with which the compound of the invention is normally associated in its natural state. Preferably, "purified," "substantially purified," and "isolated" means that the composition comprises at least 0.5%, 1%, 5%, 10%, or 20%, and most preferably at least 50% or 75% of the mass, by weight, of a given sample. In one preferred embodiment, these terms refer to the compound of the invention comprising at least 95% of the mass, by weight, of a given sample. As used herein, the terms "purified," "substantially purified," and "isolated" "isolated," when referring to a nucleic acid or protein, of nucleic acids or proteins, also refers to a state of purification or concentration different than that which occurs naturally in the mammalian, especially human, body. Any degree of purification or concentration greater than that which occurs naturally in the mammalian, especially human, body, including (1) the purification from other associated structures or compounds or (2) the association with structures or compounds to which it is not normally associated in the mammalian, especially human, body, are within the meaning of "isolated." The nucleic acid or protein or classes of nucleic acids or proteins, described herein, may be isolated, or otherwise associated with structures or compounds to which they are not normally associated in nature, according to a variety of methods and processes known to those of skill in the art.

As used herein, the term "isolated," when referring to a nucleic acid or polypeptide refers to a state of purification or concentration different than that which

occurs naturally in the mammalian, especially human, body. Any degree of purification or concentration greater than that which occurs naturally in the body, including (1) the purification from other naturally-occurring associated structures or compounds, or (2) the association with structures or compounds to which it is not normally associated in the body are within the meaning of "isolated" as used herein. The nucleic acids or polypeptides described herein may be isolated or otherwise associated with structures or compounds to which they are not normally associated in nature, according to a variety of methods and processed known to those of skill in the art.

As used herein, the terms "amplifying" and "amplification" refer to the use of any suitable amplification methodology for generating or detecting recombinant or naturally expressed nucleic acid, as described in detail, below. For example, the invention provides methods and reagents (*e.g.*, specific oligonucleotide primer pairs) for amplifying (*e.g.*, by polymerase chain reaction, PCR) naturally expressed (*e.g.*, genomic or mRNA) or recombinant (*e.g.*, cDNA) nucleic acids of the invention (*e.g.*, tastant-binding sequences of the invention) *in vivo* or *in vitro*.

The term "expression vector" refers to any recombinant expression system for the purpose of expressing a nucleic acid sequence of the invention *in vitro* or *in vivo*, constitutively or inducibly, in any cell, including prokaryotic, yeast, fungal, plant, insect or mammalian cell. The term includes linear or circular expression systems. The term includes expression systems that remain episomal or integrate into the host cell genome. The expression systems can have the ability to self-replicate or not, *i.e.*, drive only transient expression in a cell. The term includes recombinant expression "cassettes which contain only the minimum elements needed for transcription of the recombinant nucleic acid.

The term "library" means a preparation that is a mixture of different nucleic acid or poly-peptide molecules, such as the library of recombinantly generated sensory, particularly taste receptor ligand-binding regions generated by amplification of nucleic acid with degenerate primer pairs, or an isolated collection of vectors that incorporate the amplified ligand-binding regions, or a mixture of cells each randomly transfected with at least one vector encoding an taste receptor.

The term "nucleic acid" or "nucleic acid sequence" refers to a deoxy-ribonucleotide or ribonucleotide oligonucleotide in either single- or double-stranded form. The term encompasses nucleic acids, *i.e.*, oligonucleotides, containing known analogs of natural nucleotides. The term also encompasses nucleic-acid-like structures with synthetic backbones.

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating, *e.g.*, sequences in which the third position of one or more selected codons is substituted with mixed-base and/or deoxyinosine residues (Batzner *et al.*, *Nucleic Acid Res.*, 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.*, 260:2605-08 (1985); Rossolini *et al.*, *Mol. Cell. Probes*, 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

The terms "polypeptide," "peptide" and "protein" are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer.

The "translocation domain," "ligand-binding region", and chimeric receptors compositions described herein also include "analogs," or "conservative variants" and "mimetics" ("peptidomimetics") with structures and activity that substantially correspond to the exemplary sequences. Thus, the terms "conservative variant" or "analog" or "mimetic" refer to a polypeptide which has a modified amino acid sequence, such that the change(s) do not substantially alter the polypeptide's (the conservative variant's) structure and/or activity, as defined herein. These include conservatively modified variations of an amino acid sequence, *i.e.*, amino acid substitutions, additions or deletions of those residues that are not critical for protein activity, or substitution of amino acids with residues having similar properties (*e.g.*, acidic, basic, positively or negatively charged, polar or non-polar, *etc.*) such that the substitutions of even critical amino acids does not substantially alter structure and/or activity.

More particularly, "conservatively modified variants" applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not
5 encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein.

For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the
10 codon can be altered to any of the corresponding codons described without altering the encoded polypeptide.

Such nucleic acid variations are "silent variations," which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic
15 acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

20 Conservative substitution tables providing functionally similar amino acids are well known in the art. For example, one exemplary guideline to select conservative substitutions includes (original residue followed by exemplary substitution): ala/gly or ser; arg/lys; asn/gln or his; asp/glu; cys/ser; gln/asn; gly/asp; gly/ala or pro; his/asn or
25 gln; ile/leu or val; leu/ile or val; lys/arg or gln or glu; met/leu or tyr or ile; phe/met or leu or tyr; ser/thr; thr/ser; trp/tyr; tyr/trp or phe; val/ile or leu. An alternative exemplary guideline uses the following six groups, each containing amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (I); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
30 and 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); (*see also, e.g., Creighton, Proteins*, W.H. Freeman and Company (1984); Schultz and Schimer, *Principles of Protein Structure*, Springer-Verlag (1979)). One of skill in the art will appreciate that

the above-identified substitutions are not the only possible conservative substitutions. For example, for some purposes, one may regard all charged amino acids as conservative substitutions for each other whether they are positive or negative. In addition, individual substitutions, deletions or additions that alter, add or delete a
5 single amino acid or a small percentage of amino acids in an encoded sequence can also be considered "conservatively modified variations."

The terms "mimetic" and "peptidomimetic" refer to a synthetic chemical compound that has substantially the same structural and/or functional characteristics of the polypeptides, *e.g.*, translocation domains, ligand-binding regions, or chimeric
10 receptors of the invention. The mimetic can be either entirely composed of synthetic, non-natural analogs of amino acids, or may be a chimeric molecule of partly natural peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetic's structure and/or
15 activity.

As with polypeptides of the invention which are conservative variants, routine experimentation will determine whether a mimetic is within the scope of the invention, *i.e.*, that its structure and/or function is not substantially altered. Polypeptide mimetic compositions can contain any combination of non-natural
20 structural components, which are typically from three structural groups: a) residue linkage groups other than the natural amide bond ("peptide bond") linkages; b) non-natural residues in place of naturally occurring amino acid residues; or c) residues which induce secondary structural mimicry, *i.e.*, to induce or stabilize a secondary structure, *e.g.*, a beta turn, gamma turn, beta sheet, alpha helix conformation, and the
25 like. A polypeptide can be characterized as a mimetic when all or some of its residues are joined by chemical means other than natural peptide bonds. Individual peptidomimetic residues can be joined by peptide bonds, other chemical bonds or coupling means, such as, *e.g.*, glutaraldehyde, N-hydroxysuccinimide esters, bifunctional maleimides, N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-
30 diisopropylcarbodiimide (DIC). Linking groups that can be an alternative to the traditional amide bond ("peptide bond") linkages include, *e.g.*, ketomethylene (*e.g.*, -C(=O)-CH₂- for -C(=O)-NH-), aminomethylene (CH₂-NH), ethylene, olefin

(CH=CH), ether (CH₂-O), thioether (CH₂-S), tetrazole (CN₄), thiazole, retroamide, thioamide, or ester (*see, e.g.,* Spatola, *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, Vol. 7, 267-357, Marcell Dekker, *Peptide Backbone Modifications*, NY (1983)). A polypeptide can also be characterized as a mimetic by
5 containing all or some non-natural residues in place of naturally occurring amino acid residues; non-natural residues are well described in the scientific and patent literature.

A "label" or a "detectable moiety" is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ³²P, fluorescent dyes, electron-dense reagents, enzymes
10 (*e.g.,* as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins which can be made detectable, *e.g.,* by incorporating a radiolabel into the peptide or used to detect antibodies specifically reactive with the peptide.

A "labeled nucleic acid probe or oligonucleotide" is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van
15 der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe.

As used herein a "nucleic acid probe or oligonucleotide" is defined as a nucleic acid capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base
20 pairing, usually through hydrogen bond formation. As used herein, a probe may include natural (*i.e.,* A, G, C, or T) or modified bases (7-deazaguanosine, inosine, *etc.*). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester bond, so long as it does not interfere with hybridization. Thus, for example, probes may be peptide nucleic acids in which the constituent bases are
25 joined by peptide bonds rather than phosphodiester linkages. It will be understood by one of skill in the art that probes may bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are optionally directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to
30 which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or subsequence.

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, *e.g.*, a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (*e.g.*, a fusion protein).

A "promoter" is defined as an array of nucleic acid sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

As used herein, "recombinant" refers to a polynucleotide synthesized or otherwise manipulated *in vitro* (*e.g.*, "recombinant polynucleotide"), to methods of using recombinant polynucleotides to produce gene products in cells or other biological systems, or to a polypeptide ("recombinant protein") encoded by a recombinant polynucleotide. "Recombinant means" also encompass the ligation of nucleic acids having various coding regions or domains or promoter sequences from different sources into an expression cassette or vector for expression of, *e.g.*, inducible or constitutive expression of a fusion protein comprising a translocation domain of the invention and a nucleic acid sequence amplified using a primer of the invention.

The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under

stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

The phrase "stringent hybridization conditions" refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology - Hybridisation with Nucleic Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10° C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m, 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, optionally 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5x SSC, and 1% SDS, incubating at 42°C, or, 5x SSC, 1% SDS, incubating at 65°C, with wash in 0.2x SSC, and 0.1 % SDS at 65°C. Such hybridizations and wash steps can be carried out for, e.g., 1, 2, 5, 10, 15, 30, 60; or more minutes.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially related if the polypeptides which they encode are substantially related. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions.

Exemplary "moderately stringent hybridization conditions" include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 1x SSC at 45°C. Such hybridizations and wash steps can be carried out for, *e.g.*, 1, 2, 5, 10, 15, 30, 60, or more minutes. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

"Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50-70 kDa). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively.

A "chimeric antibody" is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, *e.g.*, an enzyme, toxin, hormone, growth factor, drug, *etc.*; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

An "anti-T2R" antibody is an antibody or antibody fragment that specifically binds a polypeptide encoded by a T2R gene, cDNA, or a subsequence thereof.

The term "immunoassay" is an assay that uses an antibody to specifically bind an antigen. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

The phrase “specifically (or selectively) binds” to an antibody or, “specifically (or selectively) immunoreactive with,” when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein.

For example, polyclonal antibodies raised to a T2R family member from specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with the T2R polypeptide or an immunogenic portion thereof and not with other proteins, except for orthologs or polymorphic variants and alleles of the T2R polypeptide. This selection may be achieved by subtracting out antibodies that cross-react with T2R molecules from other species or other T2R molecules. Antibodies can also be selected that recognize only T2R GPCR family members but not GPCRs from other families. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (*see, e.g., Harlow & Lane, Antibodies, A Laboratory Manual, (1988), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity*). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

The phrase “selectively associates with” refers to the ability of a nucleic acid to “selectively hybridize” with another as defined above, or the ability of an antibody to “selectively (or specifically) bind to a protein, as defined above.

The term “expression vector” refers to any recombinant expression system for the purpose of expressing a nucleic acid sequence of the invention *in vitro* or *in vivo*, constitutively or inducibly, in any cell, including prokaryotic, yeast, fungal, plant, insect or mammalian cell. The term includes linear or circular expression systems. The term includes expression systems that remain episomal or integrate into the host

cell genome. The expression systems can have the ability to self-replicate or not, *i.e.*, drive only transient expression in a cell. The term includes recombinant expression "cassettes which contain only the minimum elements needed for transcription of the recombinant nucleic acid.

5 By "host cell" is meant a cell that contains an expression vector and supports the replication or expression of the expression vector. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells such as CHO, HeLa, HEK-293, and the like, *e.g.*, cultured cells, explants, and cells *in vivo*.

10

C. Isolation and Expression of T2Rs

Isolation and expression of the T2Rs, or fragments or variants thereof, of the invention can be effected by well established cloning procedures using probes or primers constructed based on the T2R nucleic acids sequences disclosed in the application. Related T2R sequences may also be identified from human or other species genomic databases using the sequences disclosed herein and known computer-based search technologies, *e.g.*, BLAST sequence searching. In a particular embodiment, the pseudogenes disclosed herein can be used to identify functional alleles or related genes.

20 Expression vectors can then be used to infect or transfect host cells for the functional expression of these sequences. These genes and vectors can be made and expressed *in vitro* or *in vivo*. One of skill will recognize that desired phenotypes for altering and controlling nucleic acid expression can be obtained by modulating the expression or activity of the genes and nucleic acids (*e.g.*, promoters, enhancers and the like) within the vectors of the invention. Any of the known methods described for increasing or decreasing expression or activity can be used. The invention can be practiced in conjunction with any method or protocol known in the art, which are well described in the scientific and patent literature.

Alternatively, these nucleic acids can be synthesized *in vitro* by well-known chemical synthesis techniques, as described in, *e.g.*, Carruthers, *Cold Spring Harbor Symp. Quant. Biol.* 47:411-18 (1982); Adams, *Am. Chem. Soc.*, 105:661 (1983); Belousov, *Nucleic Acids Res.* 25:3440-3444 (1997); Frenkel, *Free Radic. Biol. Med.*

30

19:373-380 (1995); Blommers, *Biochemistry* 33:7886-7896 (1994); Narang, *Meth. Enzymol.* 68:90 (1979); Brown, *Meth. Enzymol.* 68:109 (1979); Beaucage, *Tetra. Lett.* 22:1859 (1981); U.S. Patent No. 4,458,066. Double-stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the
5 strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence..

Techniques for the manipulation of nucleic acids, such as, for example, for generating mutations in sequences, subcloning, labeling probes, sequencing, hybridization and the like are well described in the scientific and patent literature.
10 See, e.g., Sambrook, ed., *Molecular Cloning: A Laboratory Manual* (2nd ed.), Vols. 1-3, Cold Spring Harbor Laboratory (1989); Ausubel, ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., New York (1997); Tijssen, ed., *Laboratory Techniques in Biochemistry and Molecular Biology: Hybridization With Nucleic Acid Probes, Part I, Theory and Nucleic Acid Preparation*, Elsevier, N.Y.
15 (1993).

Nucleic acids, vectors, capsids, polypeptides, and the like can be analyzed and quantified by any of a number of general means well known to those of skill in the art. These include, e.g., analytical biochemical methods such as NMR, spectrophotometry, radiography, electrophoresis, capillary electrophoresis, high performance liquid
20 chromatography (HPLC), thin layer chromatography (TLC), and hyperdiffusion chromatography, various immunological methods, e.g., fluid or gel precipitin reactions, immunodiffusion, immunoelectrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immuno-fluorescent assays, Southern analysis, Northern analysis, dot-blot analysis, gel electrophoresis (e.g., SDS-
25 PAGE), RT-PCR, quantitative PCR, other nucleic acid or target or signal amplification methods, radiolabeling, scintillation counting, and affinity chromatography.

Oligonucleotide primers may be used to amplify nucleic acids encoding a T2R ligand-binding region. The nucleic acids described herein can also be cloned or
30 measured quantitatively using amplification techniques. Amplification methods are also well known in the art, and include, e.g., polymerase chain reaction (PCR) (Innis ed., *PCR Protocols, a Guide to Methods and Applications*, Academic Press, N.Y.

(1990); Innis ed., *PCR Strategies*, Academic Press, Inc., N.Y. (1995)); ligase chain reaction (LCR) (Wu, *Genomics*, 4:560 (1989); Landegren, *Science*, 241:1077 (1988); Barringer, *Gene*, 89:117 (1990)); transcription amplification (Kwoh, *PNAS*, 86:1173 (1989)); self-sustained sequence replication (Guatelli, *PNAS*, 87:1874 (1990)); Q Beta replicase amplification (Smith, *J. Clin. Microbiol.*, 35:1477-91 (1997)); automated Q-beta replicase amplification assay (Burg, *Mol. Cell. Probes*, 10:257-71 (1996)); and other RNA polymerase mediated techniques (e.g., NASBA, Cangene, Mississauga, Ontario). See also, Berger, *Methods Enzymol.*, 152:307-16 (1987); Sambrook; Ausubel; U.S. Patent Nos. 4,683,195 and 4,683,202; Sooknanan, *Biotechnology*, 13:563-64 (1995).

Once amplified, the nucleic acids, either individually or as libraries, may be cloned according to methods known in the art, if desired, into any of a variety of vectors using routine molecular biological methods; methods for cloning *in vitro* amplified nucleic acids are described, e.g., U.S. Pat. No. 5,426,039. To facilitate cloning of amplified sequences, restriction enzyme sites can be "built into" the PCR primer pair. For example, Pst I and Bsp E1 sites were designed into the exemplary primer pairs of the invention. These particular restriction sites have a sequence that, when ligated, are "in-frame" with respect to the 7-membrane receptor "donor" coding sequence into which they are spliced (the ligand-binding region coding sequence is internal to the 7-membrane polypeptide, thus, if it is desired that the construct be translated downstream of a restriction enzyme splice site, out of frame results should be avoided; this may not be necessary if the inserted ligand-binding region comprises substantially most of the transmembrane VII region). The primers can be designed to retain the original sequence of the "donor" 7-membrane receptor. Alternatively, the primers can encode amino acid residues that are conservative substitutions (e.g., hydrophobic for hydrophobic residue, see above discussion) or functionally benign substitutions (e.g., do not prevent plasma membrane insertion, cause cleavage by peptidase, cause abnormal folding of receptor, and the like).

The primer pairs may be designed to selectively amplify ligand-binding regions of T2R proteins. These binding regions may vary for different ligands; thus, what may be a minimal binding region for one ligand, may be too limiting for a second potential ligand. Thus, binding regions of different sizes comprising different

domain structures may be amplified; for example, transmembrane (TM) domains II through VII, III through VII, III through VI or II through VI, or variations thereof (*e.g.*, only a subsequence of a particular domain, mixing the order of the domains, and the like), of a 7-transmembrane T2R.

5 As domain structures and sequence of many 7-membrane T2R proteins are known, the skilled artisan can readily select domain-flanking and internal domain sequences as model sequences to design degenerate amplification primer pairs. For example, a nucleic acid sequence encoding domain regions II through VII can be generated by PCR amplification using a primer pair. To amplify a nucleic acid
10 comprising transmembrane domain I (TM I) sequence, a degenerate primer can be designed from a nucleic acid that encodes the amino acid sequence of the T2R family consensus sequence 1 described above. Such a degenerate primer can be used to generate a binding region incorporating TM I through TM III, TM I through TM IV, TM I through TM V, TM I through TM VI or TM I through TM VII). Other
15 degenerate primers can be designed based on the other T2R family consensus sequences provided herein. Such a degenerate primer can be used to generate a binding region incorporating TM III through TM IV, TM III through TM V, TM III through TM VI or TM III through TM VII.

Paradigms to design degenerate primer pairs are well known in the art. For
20 example, a COnsensus-DEgenerate Hybrid Oligonucleotide Primer (CODEHOP) strategy computer program is accessible as <http://blocks.fhcrc.org/codehop.html>, and is directly linked from the BlockMaker multiple sequence alignment site for hybrid primer prediction beginning with a set of related protein sequences, as known taste receptor ligand-binding regions (*see, e.g., Rose, Nucleic Acids Res.*, 26:1628-35
25 (1998); Singh, *Biotechniques*, 24:318-19 (1998)).

Means to synthesize oligonucleotide primer pairs are well known in the art. "Natural" base pairs or synthetic base pairs can be used. For example, use of artificial nucleobases offers a versatile approach to manipulate primer sequence and generate a more complex mixture of amplification products. Various families of artificial
30 nucleobases are capable of assuming multiple hydrogen bonding orientations through internal bond rotations to provide a means for degenerate molecular recognition. Incorporation of these analogs into a single position of a PCR primer allows for

generation of a complex library of amplification products. *See, e.g.,* Hoops, *Nucleic Acids Res.*, 25:4866-71 (1997). Nonpolar molecules can also be used to mimic the shape of natural DNA bases. A non-hydrogen-bonding shape mimic for adenine can replicate efficiently and selectively against a nonpolar shape mimic for thymine (*see, e.g.,* Morales, *Nat. Struct. Biol.*, 5:950-54 (1998)). For example, two degenerate bases can be the pyrimidine base 6H, 8H-3,4-dihydropyrimido[4,5-c][1,2]oxazin-7-one or the purine base N6-methoxy-2,6-diaminopurine (*see, e.g.,* Hill, *PNAS*, 95:4258-63 (1998)). Exemplary degenerate primers of the invention incorporate the nucleobase analog 5'-Dimethoxytrityl-N-benzoyl-2'-deoxy-Cytidine,3'-[(2-cyanoethyl)-(N,N-diisopropyl)]-phosphoramidite (the term "P" in the sequences, *see above*). This pyrimidine analog hydrogen bonds with purines, including A and G residues.

Polymorphic variants, alleles, and interspecies homologs that are substantially identical to a taste receptor disclosed herein can be isolated using the nucleic acid probes described above. Alternatively, expression libraries can be used to clone T2R polypeptides and polymorphic variants, alleles, and interspecies homologs thereof, by detecting expressed homologs immunologically with antisera or purified antibodies made against a T2R polypeptide, which also recognize and selectively bind to the T2R homolog.

Nucleic acids that encode ligand-binding regions of taste receptors may be generated by amplification (*e.g.,* PCR) of appropriate nucleic acid sequences using appropriate (perfect or degenerate) primer pairs. The amplified nucleic acid can be genomic DNA from any cell or tissue or mRNA or cDNA derived from taste receptor-expressing cells.

In one embodiment, hybrid protein-coding sequences comprising nucleic acids encoding T2Rs fused to a translocation sequences may be constructed. Also provided are hybrid T2Rs comprising the translocation motifs and tastant-binding regions of other families of chemosensory receptors, particularly taste receptors. These nucleic acid sequences can be operably linked to transcriptional or translational control elements, *e.g.,* transcription and translation initiation sequences, promoters and enhancers, transcription and translation terminators, polyadenylation sequences, and other sequences useful for transcribing DNA into RNA. In construction of recombinant expression cassettes, vectors, and transgenics, a promoter fragment can

be employed to direct expression of the desired nucleic acid in all desired cells or tissues.

In another embodiment, fusion proteins may include C-terminal or N-terminal translocation sequences. Further, fusion proteins can comprise additional elements, e.g., for protein detection, purification, or other applications. Detection and purification facilitating domains include, e.g., metal chelating peptides such as polyhistidine tracts, histidine-tryptophan modules, or other domains that allow purification on immobilized metals; maltose binding protein; protein A domains that allow purification on immobilized immunoglobulin; or the domain utilized in the FLACS extension/affinity purification system (Immunex, Corp, Seattle WA).

The inclusion of a cleavable linker sequences such as Factor Xa (*see, e.g., Ottavi, Biochimie*, 80:289-93 (1998)), subtilisin protease recognition motif (*see, e.g., Polyak, Protein Eng.*, 10:615-19 (1997)); enterokinase (Invitrogen, San Diego, CA), and the like, between the translocation domain (for efficient plasma membrane expression) and the rest of the newly translated polypeptide may be useful to facilitate purification. For example, one construct can include a polypeptide encoding a nucleic acid sequence linked to six histidine residues followed by a thioredoxin, an enterokinase cleavage site (*see, e.g., Williams, Biochemistry*, 34:1787-97 (1995)), and an C-terminal translocation domain. The histidine residues facilitate detection and purification while the enterokinase cleavage site provides a means for purifying the desired protein(s) from the remainder of the fusion protein. Technology pertaining to vectors encoding fusion proteins and application of fusion proteins are well described in the scientific and patent literature (*see, e.g., Kroll, DNA Cell. Biol.*, 12:441-53 (1993)).

Expression vectors, either as individual expression vectors or as libraries of expression vectors, comprising the ligand-binding region encoding sequences may be introduced into a genome or into the cytoplasm or a nucleus of a cell and expressed by a variety of conventional techniques, well described in the scientific and patent literature. *See, e.g., Roberts, Nature*, 328:731 (1987); Berger *supra*; Schneider, *Protein Expr. Purif.*, 6435:10 (1995); Sambrook; Tijssen; Ausubel. Product information from manufacturers of biological reagents and experimental equipment also provide information regarding known biological methods. The vectors can be

isolated from natural sources, obtained from such sources as ATCC or GenBank libraries, or prepared by synthetic or recombinant methods.

The nucleic acids can be expressed in expression cassettes, vectors or viruses which are stably or transiently expressed in cells (*e.g.*, episomal expression systems).

- 5 Selection markers can be incorporated into expression cassettes and vectors to confer a selectable phenotype on transformed cells and sequences. For example, selection markers can code for episomal maintenance and replication such that integration into the host genome is not required. For example, the marker may encode antibiotic resistance (*e.g.*, chloramphenicol, kanamycin, G418, bleomycin, hygromycin) or
10 herbicide resistance (*e.g.*, chlorosulfuron or Basta) to permit selection of those cells transformed with the desired DNA sequences (*see, e.g.*, Blondelet-Rouault, *Gene*, 190:315-17 (1997); Aubrecht, *J. Pharmacol. Exp. Ther.*, 281:992-97 (1997)). Because selectable marker genes conferring resistance to substrates like neomycin or hygromycin can only be utilized in tissue culture, chemoresistance genes are also used
15 as selectable markers *in vitro* and *in vivo*.

- A chimeric nucleic acid sequence may encode a T2R ligand-binding region within any 7-transmembrane polypeptide. Because 7-transmembrane receptor polypeptides have similar primary sequences and secondary and tertiary structures, structural domains (*e.g.*, extracellular domain, TM domains, cytoplasmic domain, *etc.*)
20 can be readily identified by sequence analysis. For example, homology modeling, Fourier analysis and helical periodicity detection can identify and characterize the seven domains with a 7-transmembrane receptor sequence. Fast Fourier Transform (FFT) algorithms can be used to assess the dominant periods that characterize profiles of the hydrophobicity and variability of analyzed sequences. Periodicity detection
25 enhancement and alpha helical periodicity index can be done as by, *e.g.*, Donnelly, *Protein Sci.*, 2:55-70 (1993). Other alignment and modeling algorithms are well known in the art (*see, e.g.*, Peitsch, *Receptors Channels*, 4:161-64 (1996); Kyte & Doolittle, *J. Mol. Biol.*, 157:105-32 (1982); Cronet, *Protein Eng.*, 6:59-64 (1993); <http://bioinfo.weizmann.ac.il/>).

- 30 The present invention also includes not only the nucleic acid molecules and polypeptides having the specified nucleic and amino acid sequences, but also fragments thereof, particularly fragments of, *e.g.*, 40, 60, 80, 100, 150, 200, or 250

nucleotides, or more, as well as polypeptide fragments of, *e.g.*, 10, 20, 30, 50, 70, 100, or 150 amino acids, or more. Optionally, the nucleic acid fragments can encode an antigenic polypeptide that is capable of binding to an antibody raised against a T2R family member. Further, a protein fragment of the invention can optionally be an
5 antigenic fragment that is capable of binding to an antibody raised against a T2R family member.

Also contemplated are chimeric proteins, comprising at least 10, 20, 30, 50, 70, 100, or 150 amino acids, or more, of one of at least one of the T2R polypeptides described herein, coupled to additional amino acids representing all or part of another
10 GPCR, preferably a member of the 7 transmembrane superfamily. These chimeras can be made from the instant receptors and another GPCR, or they can be made by combining two or more of the present receptors. In one embodiment, one portion of the chimera corresponds to, or is derived from the transmembrane domain of a T2R polypeptide of the invention. In another embodiment, one portion of the chimera
15 corresponds to, or is derived from the one or more of the transmembrane regions of a T2R polypeptide described herein, and the remaining portion or portions can come from another GPCR. Chimeric receptors are well known in the art, and the techniques for creating them and the selection and boundaries of domains or fragments of G Protein-Coupled Receptors for incorporation therein are also well known. Thus, this
20 knowledge of those skilled in the art can readily be used to create such chimeric receptors. The use of such chimeric receptors can provide, for example, a taste selectivity characteristic of one of the receptors specifically disclosed herein, coupled with the signal transduction characteristics of another receptor, such as a well known receptor used in prior art assay systems.

For example, a region such as a ligand-binding region, an extracellular domain, a transmembrane domain, a transmembrane domain, a cytoplasmic domain, an N-terminal domain, a C-terminal domain, or any combination thereof, can be covalently linked to a heterologous protein. For instance, a T2R transmembrane region can be linked to a heterologous GPCR transmembrane domain, or a
30 heterologous GPCR extracellular domain can be linked to a T2R transmembrane region. Other heterologous proteins of choice can include, *e.g.*, green fluorescent protein, β -gal, glutamate receptor, and the rhodopsin N-terminus.

Also within the scope of the invention are host cells for expressing the T2Rs, fragments, or variants of the invention. To obtain high levels of expression of a cloned gene or nucleic acid, such as cDNAs encoding the T2Rs, fragments, or variants of the invention, one of skill typically subclones the nucleic acid sequence of interest
5 into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and if for a nucleic acid encoding a protein, a ribosome binding site for translational initiation. Suitable bacterial promoters are well known in the art and described, *e.g.*, in Sambrook *et al.* However, bacterial or eukaryotic expression systems can be used.

10 Any of the well-known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a
15 host cell (*see, e.g.*, Sambrook *et al.*) It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one nucleic acid molecule into the host cell capable of expressing the T2R, fragment, or variant of interest.

After the expression vector is introduced into the cells, the transfected cells are
20 cultured under conditions favoring expression of the receptor, fragment, or variant of interest, which is then recovered from the culture using standard techniques. Examples of such techniques are well known in the art. *See, e.g.*, WO 00/06593, which is incorporated by reference in a manner consistent with this disclosure.

25 D. Immunological Detection of T2Rs

In addition to the detection of T2R genes and gene expression using nucleic acid hybridization technology, one can also use immunoassays to detect T2Rs, *e.g.*, to identify taste receptor cells, and variants of T2R family members. Immunoassays can be used to qualitatively or quantitatively analyze the T2Rs. A general overview of the
30 applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

1. Antibodies to T2R family members

Methods of producing polyclonal and monoclonal antibodies that react specifically with a T2R family member are known to those of skill in the art (*see, e.g., Coligan, Current Protocols in Immunology* (1991); Harlow & Lane, *supra*; Goding, 5 *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986); and Kohler & Milstein, *Nature*, 256:495-97 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors, as well as preparation of polyclonal and monoclonal antibodies by immunizing rabbits or mice (*see, e.g., Huse et al., Science*, 246:1275-81 (1989); Ward 10 *et al., Nature*, 341:544-46 (1989)).

A number of T2R-comprising immunogens may be used to produce antibodies specifically reactive with a T2R family member. For example, a recombinant T2R protein, or an antigenic fragment thereof, can be isolated as described herein. Suitable antigenic regions include, *e.g.*, the consensus sequences disclosed above that can be 15 used to identify members of the T2R family. Recombinant proteins can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally known in the art. Recombinant protein is a preferred immunogen for the production of monoclonal or polyclonal antibodies. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used an 20 immunogen. Naturally occurring protein may also be used either in pure or impure form. The product may then be injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated, for subsequent use in immunoassays to measure the protein.

More specifically, methods of production of polyclonal antibodies are known 25 to those of skill in the art. For example, an inbred strain of mice (*e.g.*, BALB/C mice) or rabbits may be immunized with a T2R polypeptide using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. The animal's immune response to the immunogen preparation may then be monitored by taking test bleeds and determining the titer of reactivity to the T2R. When appropriately high titers of 30 antibody to the immunogen are obtained, blood may be collected from the animal and antisera may be prepared. Further fractionation of the antisera to enrich for antibodies reactive to the T2R polypeptide can be done if desired (*see Harlow & Lane, supra*).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen may be immortalized, commonly by fusion with a myeloma cell (*see* Kohler & Milstein, *Eur. J. Immunol.*, 6:511-19 (1976)). Alternative methods of immortalization
5 include transformation with Epstein Barr Virus, oncogenes, retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells may then be screened for production of antibodies of the desired specificity and affinity for the antigen. Yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a
10 vertebrate host. Alternatively, one may isolate nucleic acid sequences that encode a monoclonal antibody, or a binding fragment thereof, by screening a nucleic acid library from human B cells according to the general protocol outlined by Huse *et al.*, *Science*, 246:1275-81 (1989).

Monoclonal antibodies and polyclonal sera are generally collected and titrated
15 against the immunogen protein in an immunoassay, *e.g.*, a solid phase immunoassay with the immunogen immobilized on a solid support. Typically, polyclonal antisera with a titer of 10⁴ or greater may be selected and tested for their cross reactivity against non-T2R proteins, or even other T2R family members or other related proteins from other organisms, using a competitive binding immunoassay. Specific polyclonal
20 antisera and monoclonal antibodies will usually bind with a K_d of at least about 0.1 mM, more usually at least about 1 pM, optionally at least about 0.1 pM or better, and optionally 0.01 pM or better.

Once T2R family member specific antibodies are available, individual T2R proteins can be detected by a variety of immunoassay methods. For a review of
25 immunological and immunoassay procedures, *see* Stites & Terr eds., *Basic and Clinical Immunology* (7th ed. 1991). Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in Maggio, ed., *Enzyme Immunoassay* (1980); and Harlow & Lane, *supra*.

2. Immunological binding assays

30 T2R proteins can be detected and/or quantified using any of a number of well recognized immunological binding assays (*see, e.g.*, U.S. Patents 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, *see*

Asai, ed., *Methods in Cell Biology: Antibodies in Cell Biology*, volume 37 (1993); Stites & Terr, eds., *Basic and Clinical Immunology* (7th ed. 1991). Immunological binding assays (or immunoassays) typically use an antibody that specifically binds to a protein or antigen of choice (in this case a T2R family member or an antigenic subsequence thereof). The antibody (e.g., anti-T2R) may be produced by any of a number of means well known to those of skill in the art, as described above.

Immunoassays also often use a labeling agent to specifically bind to, and label the complex formed by the antibody and antigen. The labeling agent may itself be one of the moieties comprising the antibody/antigen complex. Thus, the labeling agent may be a labeled T2R polypeptide or a labeled anti-T2R antibody. Alternatively, the labeling agent may be a third moiety, e.g., a secondary antibody, that specifically binds to the antibody/T2R complex (a secondary antibody is typically specific to antibodies of the species from which the first antibody is derived). Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (see, e.g., Kronval *et al.*, *J. Immunol.*, 111:1401-06 (1973); Akerstrom *et al.*, *J. Immunol.*, 135:2589-642 (1985)). The labeling agent can be modified with a detectable moiety, such as biotin, to which another molecule can specifically bind, such as streptavidin. A variety of detectable moieties are well known to those skilled in the art.

Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incubation steps can vary from about 5 seconds to several hours, optionally from about 5 minutes to about 24 hours. However, the incubation time will generally depend upon the assay format, antigen, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient temperature, although they can be conducted over a range of temperatures, e.g., from about 10°C to about 40°C.

a. Non-competitive assay formats

Immunoassays for detecting a T2R protein in a sample may be either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of antigen is directly measured. In one preferred "sandwich" assay, for

example, the anti-T2R antibodies can be bound directly to a solid substrate on which they are immobilized. These immobilized antibodies may then capture any T2R protein present in the test sample. The T2R protein is thus immobilized, and is then bound by a labeling agent, such as a second T2R antibody bearing a label.

- 5 Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second or third antibody is typically modified with a detectable moiety, such as biotin, to which another molecule specifically binds, *e.g.*, streptavidin, to provide a detectable moiety.

10 **b. Competitive assay formats**

- In competitive assays, the amount of T2R protein present in the sample is measured indirectly by measuring the amount of a known, added (exogenous) T2R protein displaced (competed away) from an anti-T2R antibody by the unknown T2R protein present in a sample. In one competitive assay, a known amount of T2R protein is added to a sample, and the sample is then contacted with an antibody that specifically binds to the T2R. The amount of exogenous T2R protein bound to the antibody is inversely proportional to the concentration of T2R protein present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of T2R protein bound to the antibody may be determined
- 15 either by measuring the amount of T2R protein present in a T2R/antibody complex, or alternatively by measuring the amount of remaining uncomplexed protein. The amount of T2R protein may be detected by providing a labeled T2R molecule.

- A hapten inhibition assay is another preferred competitive assay. In this assay the known T2R protein is immobilized on a solid substrate. A known amount of anti-T2R antibody is added to the sample, and the sample is then contacted with the immobilized T2R. The amount of anti-T2R antibody bound to the known immobilized T2R protein is inversely proportional to the amount of T2R protein present in the sample. Again, the amount of immobilized antibody may be detected by detecting either the immobilized fraction of antibody or the fraction of the antibody that remains in solution. Detection may be direct where the antibody is labeled, or indirect by the subsequent addition of a labeled moiety that specifically binds to the antibody as described above.
- 25
- 30

c. Cross-reactivity determinations

Immunoassays in the competitive binding format can also be used for cross-reactivity determinations. For example, a protein at least partially encoded by the nucleic acid sequences disclosed herein can be immobilized to a solid support.

5 Proteins (*e.g.*, T2R polypeptides and homologs thereof) may be added to the assay and thereby compete for binding of the antisera to the immobilized antigen. The ability of the added proteins to compete for binding of the antisera to the immobilized protein is compared to the ability of the T2R polypeptide encoded by the nucleic acid sequences disclosed herein to compete with itself. The percent cross-reactivity for the above

10 proteins is calculated, using standard calculations. Those antisera with less than 10% cross-reactivity with each of the added proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with the added considered proteins, *e.g.*, distantly related homologs. In addition, peptides comprising amino acid sequences representing

15 consensus sequences that may be used to identify members of the T2R family can be used in cross-reactivity determinations.

The immunoabsorbed and pooled antisera may then be used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps an allele or polymorphic variant of a T2R family member, to the immunogen

20 protein (*i.e.*, T2R protein encoded by the nucleic acid sequences disclosed herein). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required to inhibit 50% of binding is less than 10 times the amount of the

25 protein encoded by nucleic acid sequences disclosed herein required to inhibit 50% of binding, then the second protein is said to specifically bind to the polyclonal antibodies generated to a T2R immunogen.

Antibodies raised against T2R consensus sequences can also be used to prepare antibodies that specifically bind only to GPCRs of the T2R family, but not to

30 GPCRs from other families. For example, polyclonal antibodies that specifically bind to a particular member of the T2R family can be made by subtracting out cross-reactive antibodies using other T2R family members. Species-specific polyclonal

antibodies can be made in a similar way. For example, antibodies specific to human T2R1 can be made by, subtracting out antibodies that are cross-reactive with orthologous sequences, *e.g.*, rat T2R1 or mouse T2R1.

d. Other assay formats

5 Western blot (immunoblot) analysis may be used to detect and quantify the presence of T2R protein in a sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (*e.g.*, a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with antibodies that
10 specifically bind the T2R protein. The anti-T2R polypeptide antibodies then specifically bind to the T2R polypeptide on the solid support. These antibodies may be directly labeled, or alternatively may be subsequently detected using labeled antibodies (*e.g.*, labeled sheep anti-mouse antibodies) that specifically bind to the anti-T2R antibodies.

15 Other, assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (*e.g.*, antibodies) and release encapsulated reagents or markers. The released chemicals are then detected according to standard techniques (*see Monroe et al., Amer. Clin. Prod. Rev.*, 5:34-41 (1986)).

e. Reduction of non-specific binding

20 One of skill in the art will appreciate that it is often desirable to minimize non-specific binding in immunoassays. Particularly, where the assay involves an antigen or antibody immobilized on a solid substrate it is desirable to minimize the amount of non-specific binding to the substrate. Means of reducing such non-specific binding are well known to those of skill in the art. Typically, these techniques involve coating
25 the substrate with a proteinaceous composition. In particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used, with powdered milk being most preferred.

f. Labels

The particular label or detectable group used in the assay is not a critical aspect
30 of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well

developed in the field of immunoassays and, in general, most any label useful in such methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical, or chemical means. Useful labels in the present invention include magnetic
5 beads (*e.g.*, DYNABEADSTM), fluorescent dyes (*e.g.*, fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (*e.g.*, ³H, ¹²⁵I, ³⁵S, ¹⁴C, or ³²P), enzymes (*e.g.*, horseradish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic beads (*e.g.*, polystyrene, polypropylene, latex, *etc.*).

10 The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide variety of labels may be used, with the choice of label depending on sensitivity required, ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

15 Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (*e.g.*, biotin) is covalently bound to the molecule. The ligand then binds to another molecules (*e.g.*, streptavidin) molecule, which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. The ligands and their
20 targets can be used in any suitable combination with antibodies that recognize a T2R protein, or secondary antibodies that recognize anti-T2R.

The molecules can also be conjugated directly to signal generating compounds, *e.g.*, by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and
25 glycosidases, or oxidotases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, *etc.* Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, *e.g.*, luminol. For a review of various labeling or signal producing systems that may be used, *see* U.S. Patent No. 4,391,904.

30 Means of detecting labels are well known to those of skill in the art. Thus, *e.g.*, where the label is a radioactive label, means for detection include a scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent

label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge-coupled devices (CCDs) or photomultipliers and the like. Similarly, enzymatic labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple colorimetric labels may be detected simply by observing the color associated with the label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labeled and the presence of the target antibody is detected by simple visual inspection.

E. Detection of taste modulators

Methods and compositions for determining whether a test compound specifically binds to a T2R polypeptide of the invention, both *in vitro* and *in vivo* are described below. Many aspects of cell physiology can be monitored to assess the effect of ligand-binding to a naturally occurring or chimeric T2Rs. These assays may be performed on intact cells expressing a T2R polypeptide, on permeabilized cells, or on membrane fractions produced by standard methods.

Taste receptors bind tastants and initiate the transduction of chemical stimuli into electrical signals. An activated or inhibited G Protein will in turn alter the properties of target enzymes, channels, and other effector proteins. Some examples are the activation of cGMP phosphodiesterase by transducin in the visual system, adenylate cyclase by the stimulatory G Protein, phospholipase C by Gq and other cognate G Proteins, and modulation of diverse channels by Gi and other G Proteins. Downstream consequences can also be examined such as generation of diacyl glycerol and IP3 by phospholipase C, and in turn, for calcium mobilization by IP3.

The T2R proteins or polypeptides of the assay will typically be selected from a polypeptide having a sequence of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or fragments or conservatively modified variants thereof.

Alternatively, the T2R proteins or polypeptides of the assay can be derived
5 from a eukaryote host cell, and can include an amino acid subsequence having amino acid sequence identity to SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or conservatively modified variants thereof. Generally, the amino acid sequence identity will be at least 30% preferably 30-40%, more specifically 50-60, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%. Optionally, the T2R proteins or
10 polypeptides of the assays can comprise a region of a T2R polypeptide, such as an extracellular domain, transmembrane region, cytoplasmic domain, ligand-binding domain, and the like. Optionally, the T2R polypeptide, or a portion thereof, can be covalently linked to a heterologous protein to create a chimeric protein used in the assays described herein.

15 Modulators of T2R activity may be tested using T2R proteins or polypeptides as described above, either recombinant or naturally occurring. The T2R proteins or polypeptides can be isolated, expressed in a cell, expressed in a membrane derived from a cell, expressed in tissue or in an animal, either recombinant or naturally occurring. For example, tongue slices, dissociated cells from a tongue, transformed
20 cells, or membranes can be used. Modulation can be tested using one of the *in vitro* or *in vivo* assays described herein.

1. In vitro binding assays

Taste transduction can also be examined *in vitro* with soluble or solid state reactions, using a T2R polypeptide or a chimeric molecule, such as an extracellular
25 domain, transmembrane region, or combination thereof, of a T2R covalently linked to a heterologous signal transduction domain; or a heterologous extracellular domain and/or transmembrane region covalently linked to the transmembrane and/or cytoplasmic domain of a T2R protein or polypeptide. Furthermore, ligand-binding regions of a T2R polypeptide can be used *in vitro* in soluble or solid state reactions to
30 assay for ligand binding. In numerous embodiments, a chimeric receptor can be made that comprises all or part of a T2R polypeptide, as well an additional sequence that

facilitates the localization of the T2R to the membrane, such as a rhodopsin, *e.g.*, an N-terminal fragment of a rhodopsin protein.

Ligand binding to a T2R protein, a ligand-binding region, or chimeric protein can be tested in solution, in a bilayer membrane, attached to a solid phase, in a lipid
5 monolayer, or in vesicles. Binding of a modulator can be tested using, *e.g.*, changes in spectroscopic characteristics (*e.g.*, fluorescence, absorbance, refractive index) hydrodynamic (*e.g.*, shape), chromatographic, or solubility properties.

T2R-G Protein interactions can also be examined. For example, binding of the G Protein to the T2R polypeptide or its release from the polypeptide can be examined.
10 In the absence of GTP, an activator will lead to the formation of a tight complex of a G protein (all three subunits) with the T2R. This complex can be detected in a variety of ways, as noted above. Such an assay can be modified to search for inhibitors, *e.g.*, by adding an activator to the T2R and G protein in the absence of GTP, which form a tight complex, and then screen for inhibitors by looking at dissociation of the T2R-G
15 protein complex. In the presence of GTP, release of the alpha subunit of the G protein from the other two G protein subunits serves as a criterion of activation.

In another embodiment of the invention, a GTP γ S assay may be used. As described above, upon activation of a GPCR, the G α subunit of the G protein complex is stimulated to exchange bound GDP for GTP. Ligand-mediated stimulation of G
20 protein exchange activity can be measured in a biochemical assay measuring the binding of added radioactively-labeled GTP γ ³⁵S to the G protein in the presence of a putative ligand. Typically, membranes containing the chemosensory receptor of interest are mixed with a complex of G proteins. Potential inhibitors and/or activators and GTP γ S are added to the assay, and binding of GTP γ S to the G protein is
25 measured. Binding can be measured by liquid scintillation counting or by any other means known in the art, including scintillation proximity assays (SPA). In other assays formats, fluorescently-labeled GTP γ S can be utilized.

In particularly preferred embodiments, T2R-gustducin interactions may be monitored as a function of T2R receptor activation. For instance, mouse T2R5 shows
30 strong cycloheximide dependent coupling with gustducin. Such ligand dependent coupling of T2R receptors with gustducin can be used as a marker to identify modifiers of any member of the T2R family.

2. Fluorescence Polarization Assays

In another embodiment, Fluorescence Polarization ("FP") based assays may be used to detect and monitor ligand binding. Fluorescence polarization is a versatile laboratory technique for measuring equilibrium binding, nucleic acid hybridization, and enzymatic activity. Fluorescence polarization assays are homogeneous in that they do not require a separation step such as centrifugation, filtration, chromatography, precipitation, or electrophoresis. These assays are done in real time, directly in solution and do not require an immobilized phase. Polarization values can be measured repeatedly and after the addition of reagents since measuring the polarization is rapid and does not destroy the sample. Generally, this technique can be used to measure polarization values of fluorophores from low picomolar to micromolar levels. This section describes how fluorescence polarization can be used in a simple and quantitative way to measure the binding of ligands to the T2R polypeptides of the invention.

When a fluorescently labeled molecule is excited with plane polarized light, it emits light that has a degree of polarization that is inversely proportional to its molecular rotation. Large fluorescently labeled molecules remain relatively stationary during the excited state (4 nanoseconds in the case of fluorescein) and the polarization of the light remains relatively constant between excitation and emission. Small fluorescently labeled molecules rotate rapidly during the excited state and the polarization changes significantly between excitation and emission. Therefore, small molecules have low polarization values and large molecules have high polarization values. For example, a single-stranded fluorescein-labeled oligonucleotide has a relatively low polarization value but when it is hybridized to a complementary strand, it has a higher polarization value. When using FP to detect and monitor tastant-binding which may activate or inhibit the chemosensory receptors of the invention, fluorescence-labeled tastants or auto-fluorescent tastants may be used.

Fluorescence polarization (P) is defined as:

$$P = \frac{Int_{\parallel} - Int_{\perp}}{Int_{\parallel} + Int_{\perp}}$$

Where \parallel is the intensity of the emission light parallel to the excitation light plane and \perp is the intensity of the emission light perpendicular to the excitation

light plane. P, being a ratio of light intensities, is a dimensionless number. For example, the Beacon ® and Beacon 2000™ System may be used in connection with these assays. Such systems typically express polarization in millipolarization units (1 Polarization Unit = 1000 mP Units).

5 The relationship between molecular rotation and size is described by the Perrin equation. Summarily, the Perrin equation states that polarization is directly proportional to the rotational relaxation time, the time that it takes a molecule to rotate through an angle of approximately 68.5°. Rotational relaxation time is related to viscosity (η), absolute temperature (T), molecular volume (V), and the gas constant
10 (R) by the following equation:

$$\text{Rotational Relaxation Time} = \frac{3\eta V}{RT}$$

The rotational relaxation time is small (≈ 1 nanosecond) for small molecules (e.g. fluorescein) and large (≈ 100 nanoseconds) for large molecules (e.g. immunoglobulins). If viscosity and temperature are held constant, rotational
15 relaxation time, and therefore polarization, is directly related to the molecular volume. Changes in molecular volume may be due to interactions with other molecules, dissociation, polymerization, degradation, hybridization, or conformational changes of the fluorescently labeled molecule. For example, fluorescence polarization has been used to measure enzymatic cleavage of large fluorescein labeled polymers by
20 proteases, DNases, and RNases. It also has been used to measure equilibrium binding for protein/protein interactions, antibody/antigen binding, and protein/DNA binding.

3. Solid state and soluble high throughput assays

In yet another embodiment, the invention provides soluble assays using a T2R polypeptide; or a cell or tissue expressing a T2R polypeptide. In another embodiment,
25 the invention provides solid phase based *in vitro* assays in a high throughput format, where the T2R polypeptide, or cell or tissue expressing the T2R polypeptide is attached to a solid phase substrate.

In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each
30 well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed,

every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (*e.g.*, 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 1000 to about 1500 different compounds. It is also possible to assay multiple compounds in each plate well. Further, it is possible to
5 assay several different plates per day allowing for assay screens of about 6,000-20,000 compounds per day. More recently, microfluidic approaches to reagent manipulation have been developed.

The molecule of interest can be bound to the solid state component, directly or indirectly, via covalent or non-covalent linkage, *e.g.*, via a tag. The tag can be any of
10 a variety of components. In general, a molecule which binds the tag (a tag binder) is fixed to a solid support, and the tagged molecule of interest (*e.g.*, the taste transduction molecule of interest) is attached to the solid support by interaction of the tag and the tag binder.

A number of tags and tag binders can be used, based upon known molecular
15 interactions well described in the literature. For example, where a tag has a natural binder, for example, biotin, protein A, or protein G, it can be used in conjunction with appropriate tag binders (avidin, streptavidin, neutravidin, the Fc region of an immunoglobulin, *etc.*). Antibodies to molecules with natural binders such as biotin are also widely available and appropriate tag binders (*see*, SIGMA Immunochemicals
20 1998 catalogue, SIGMA, St. Louis MO).

Similarly, any haptenic or antigenic compound can be used in combination with an appropriate antibody to form a tag/tag binder pair. Thousands of specific antibodies are commercially available and many additional antibodies are described in the literature. For example, in one common configuration, the tag is a first antibody
25 and the tag binder is a second antibody which recognizes the first antibody. In addition to antibody-antigen interactions, receptor-ligand interactions are also appropriate as tag and tag-binder pairs. For example, agonists and antagonists of cell membrane receptors (*e.g.*, cell receptor-ligand interactions such as transferrin, c-kit, viral receptor ligands, cytokine receptors, chemokine receptors, interleukin receptors,
30 immunoglobulin receptors and antibodies, the cadherein family, the integrin family, the selectin family, and the like; *see, e.g.*, Pigott & Power, *The Adhesion Molecule Facts Book I* (1993)). Similarly, toxins and venoms, viral epitopes, hormones (*e.g.*,

opiates, steroids, *etc.*), intracellular receptors (*e.g.*, which mediate the effects of various small ligands, including steroids, thyroid hormone, retinoids and vitamin D; peptides), drugs, lectins, sugars, nucleic acids (both linear and cyclic polymer configurations), oligosaccharides, proteins, phospholipids and antibodies can all
5 interact with various cell receptors.

Synthetic polymers, *e.g.*, polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, and polyacetates, can also form an appropriate tag or tag binder. Many other tag/tag binder pairs are also useful in assay systems described herein, as would be apparent to
10 one of skill upon review of this disclosure.

Common linkers such as peptides, polyethers, and the like can also serve as tags, and include polypeptide sequences, such as poly gly sequences of between about 5 and 200 amino acids. Such flexible linkers are known to persons of skill in the art. For example, poly(ethylene glycol) linkers are available from Shearwater Polymers,
15 Inc. Huntsville, Alabama. These linkers optionally have amide linkages, sulfhydryl linkages, or heterofunctional linkages.

Tag binders may be fixed to solid substrates using any of a variety of methods currently available. Solid substrates are commonly derivatized or functionalized by exposing all or a portion of the substrate to a chemical reagent which fixes a chemical
20 group to the surface that is reactive with a portion of the tag binder. For example, groups that are suitable for attachment to a longer chain portion include amines, hydroxyl, thiol, and carboxyl groups. Aminoalkylsilanes and hydroxyalkylsilanes can be used to functionalize a variety of surfaces, such as glass surfaces. The construction of such solid phase biopolymer arrays is well described in the literature. *See, e.g.*,
25 Merrifield, *J. Am. Chem. Soc.*, 85:2149-54 (1963) (describing solid phase synthesis of, *e.g.*, peptides); Geysen *et al.*, *J. Immun. Meth.*, 102:259-74 (1987) (describing synthesis of solid phase components on pins); Frank & Doring, *Tetrahedron*, 44:60316040 (1988) (describing synthesis of various peptide sequences on cellulose disks); Fodor *et al.*, *Science*, 251:767-77 (1991); Sheldon *et al.*, *Clinical Chemistry*,
30 39(4):718-19 (1993); and Kozal *et al.*, *Nature Medicine*, 2(7):753759 (1996) (all describing arrays of biopolymers fixed to solid substrates). Non-chemical approaches

for fixing tag binders to substrates include other common methods, such as heat, cross-linking by UV radiation, and the like.

4. Computer-based assays

Yet another assay for compounds that modulate T2R polypeptide activity involves computer assisted compound design, in which a computer system is used to generate a three-dimensional structure of an T2R polypeptide based on the structural information encoded by its amino acid sequence. The input amino acid sequence interacts directly and actively with a preestablished algorithm in a computer program to yield secondary, tertiary, and quaternary structural models of the protein. The models of the protein structure are then examined to identify regions of the structure that have the ability to bind, *e.g.*, ligands. These regions are then used to identify ligands that bind to the protein.

The three-dimensional structural model of the protein is generated by entering protein amino acid sequences of at least 10 amino acid residues or corresponding nucleic acid sequences encoding a T2R polypeptide into the computer system. The nucleotide sequence encoding the T2R polypeptide, or the amino acid sequence thereof, can be any sequence disclosed herein, and conservatively modified versions thereof.

The amino acid sequence represents the primary sequence or subsequence of the protein, which encodes the structural information of the protein. At least 10 residues of the amino acid sequence (or a nucleotide sequence encoding 10 amino acids) are entered into the computer system from computer keyboards, computer readable substrates that include, but are not limited to, electronic storage media (*e.g.*, magnetic diskettes, tapes, cartridges, and chips), optical media (*e.g.*, CD ROM), information distributed by internet sites, and by RAM. The three-dimensional structural model of the protein is then generated by the interaction of the amino acid sequence and the computer system, using software known to those of skill in the art.

The amino acid sequence represents a primary structure that encodes the information necessary to form the secondary, tertiary and quaternary structure of the protein of interest. The software looks at certain parameters encoded by the primary sequence to generate the structural model. These parameters are referred to as "energy terms," and primarily include electrostatic potentials, hydrophobic potentials, solvent

accessible surfaces, and hydrogen bonding. Secondary energy terms include van der Waals potentials. Biological molecules form the structures that minimize the energy terms in a cumulative fashion. The computer program is therefore using these terms encoded by the primary structure or amino acid sequence to create the secondary structural model.

The tertiary structure of the protein encoded by the secondary structure is then formed on the basis of the energy terms of the secondary structure. The user at this point can enter additional variables such as whether the protein is membrane bound or soluble, its location in the body, and its cellular location, *e.g.*, cytoplasmic, surface, or nuclear. These variables along with the energy terms of the secondary structure are used to form the model of the tertiary structure. In modeling the tertiary structure, the computer program matches hydrophobic faces of secondary structure with like, and hydrophilic faces of secondary structure with like.

Once the structure has been generated, potential ligand-binding regions are identified by the computer system. Three-dimensional structures for potential ligands are generated by entering amino acid or nucleotide sequences or chemical formulas of compounds, as described above. The three-dimensional structure of the potential ligand is then compared to that of the T2R polypeptide to identify ligands that bind to the protein. Binding affinity between the protein and ligands is determined using energy terms to determine which ligands have an enhanced probability of binding to the protein.

Computer systems are also used to screen for mutations, polymorphic variants, alleles, and interspecies homologs of T2R genes. Such mutations can be associated with disease states or genetic traits. As described above, GeneChip™ and related technology can also be used to screen for mutations, polymorphic variants, alleles, and interspecies homologs. Once the variants are identified, diagnostic assays can be used to identify patients having such mutated genes. Identification of the mutated T2R genes involves receiving input of a first nucleic acid or amino acid sequence of a T2R gene, or conservatively modified versions thereof. The sequence is entered into the computer system as described above. The first nucleic acid or amino acid sequence is then compared to a second nucleic acid or amino acid sequence that has substantial identity to the first sequence. The second sequence is entered into the computer

system in the manner described above. Once the first and second sequences are compared, nucleotide or amino acid differences between the sequences are identified. Such sequences can represent allelic differences in various T2R genes, and mutations associated with disease states and genetic traits.

5 **5. Cell-based binding assays**

In a preferred embodiment, a T2R protein or polypeptide is expressed in a eukaryotic cell as a chimeric receptor with a heterologous, chaperone sequence that facilitates its maturation and targeting through the secretory pathway. In a preferred embodiment, the heterologous sequence is a rhodopsin sequence, such as an N-
10 terminal fragment of a rhodopsin. Such chimeric T2R proteins can be expressed in any eukaryotic cell, such as HEK-293 cells. Preferably, the cells comprise a functional G Protein, *e.g.*, G α 15, that is capable of coupling the chimeric receptor to an intracellular signaling pathway or to a signaling protein such as phospholipase C. Activation of such chimeric receptors in such cells can be detected using any standard
15 method, such as by detecting changes in intracellular calcium by detecting FURA-2 dependent fluorescence in the cell.

Activated GPCR receptors become substrates for kinases that phosphorylate the C-terminal tail of the receptor (and possibly other sites as well). Thus, activators will promote the transfer of ^{32}P from gamma-labeled GTP to the receptor, which can
20 be assayed with a scintillation counter. The phosphorylation of the C-terminal tail will promote the binding of arrestin-like proteins, and will interfere with the binding of G Proteins. The kinase/arrestin pathway plays a key role in the desensitization of many GPCRs. For example, compounds that modulate the duration a taste receptor stays active may be useful as a means of prolonging a desired taste or cutting off an
25 unpleasant one. For a general review of GPCR signal transduction and methods of assaying signal transduction, *see, e.g., Methods in Enzymology*, vols. 237 and 238 (1994) and volume 96 (1983); Bourne *et al.*, *Nature*, 10:349:117-27 (1991); Bourne *et al.*, *Nature*, 348:125-32 (1990); Pitcher *et al.*, *Annu. Rev. Biochem.*, 67:653-92 (1998).

T2R modulation may be assayed by comparing the response of a T2R
30 polypeptide treated with a putative T2R modulator to the response of an untreated control sample. Such putative T2R modulators can include tastants that either inhibit or activate T2R polypeptide activity. In one embodiment, control samples (untreated

with activators or inhibitors) are assigned a relative T2R activity value of 100. Inhibition of a T2R polypeptide is achieved when the T2R activity value relative to the control is about 90%, optionally 50% or 25-0%. Activation of a T2R polypeptide is achieved when the T2R activity value relative to the control is 110%, optionally
5 150%, 200-500%, or 1000-2000%.

Changes in ion flux may be assessed by determining changes in ionic polarization (*i.e.*, electrical potential) of the cell or membrane expressing a T2R polypeptide. One means to determine changes in cellular polarization is by measuring changes in current (thereby measuring changes in polarization) with voltage-clamp
10 and patch-clamp techniques (*see, e.g.*, the “cell-attached” mode, the “inside-out” mode, and the “whole cell” mode, *e.g.*, Ackerman *et al.*, *New Engl. J Med.*, 336:1575-95 (1997)). Whole cell currents are conveniently determined using known standards. Other known assays include: radiolabeled ion flux assays and fluorescence assays using voltage-sensitive dyes (*see, e.g.*, Vestergaard-Bogind *et al.*, *J. Membrane Biol.*,
15 88:67-75 (1988); Gonzales & Tsien, *Chem. Biol.*, 4:269-77 (1997); Daniel *et al.*, *J. Pharmacol. Meth.*, 25:185-93 (1991); Holevinsky *et al.*, *J. Membrane Biology*, 137:59-70 (1994)). Generally, the compounds to be tested are present in the range from 1 pM to 100 mM.

The effects of test compounds upon the function of the T2R polypeptides can
20 be measured by examining any of the parameters described above. Any suitable physiological change that affects GPCR activity can be used to assess the influence of a test compound on the T2R polypeptides of this invention. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as transmitter release, hormone release, transcriptional changes
25 to both known and uncharacterized genetic markers (*e.g.*, northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as Ca^{2+} , IP3, cGMP, or cAMP.

Preferred assays for GPCRs include cells that are loaded with ion or voltage sensitive dyes to report receptor activity. Assays for determining activity of such
30 receptors can also use known agonists and antagonists for other G Protein-Coupled Receptors as negative or positive controls to assess activity of tested compounds. In assays for identifying modulatory compounds (*e.g.*, agonists, antagonists), changes in

the level of ions in the cytoplasm or membrane voltage may be monitored using an ion sensitive or membrane voltage fluorescent indicator, respectively. Among the ion-sensitive indicators and voltage probes that may be employed are those disclosed in the Molecular Probes 1997 Catalog. For GPCRs, promiscuous G proteins such as

5 G α 15 and G α 16 can be used in the assay of choice (Wilkie *et al*, *PNAS*, 88:10049-53 (1991)). Such promiscuous G proteins allow coupling of a wide range of receptors.

Receptor activation typically initiates subsequent intracellular events, *e.g.*, increases in second messengers such as IP₃, which releases intracellular stores of calcium ions. Activation of some GPCRs stimulates the formation of inositol

10 triphosphate (IP₃) through phospholipase C-mediated hydrolysis of phosphatidylinositol (Berridge & Irvine, *Nature*, 312:315-21 (1984)). IP₃ in turn stimulates the release of intracellular calcium ion stores. Thus, a change in cytoplasmic calcium ion levels, or a change in second messenger levels such as IP₃ can be used to assess GPCR function. Cells expressing such GPCRs may exhibit

15 increased cytoplasmic calcium levels as a result of contribution from both intracellular stores and via activation of ion channels, in which case it may be desirable although not necessary to conduct such assays in calcium-free buffer, optionally supplemented with a chelating agent such as EGTA, to distinguish fluorescence response resulting from calcium release from internal stores.

Other assays can involve determining the activity of receptors which, when

20 activated, result in a change in the level of intracellular cyclic nucleotides, *e.g.*, cAMP or cGMP, by activating or inhibiting enzymes such as adenylate cyclase. There are cyclic nucleotide-gated ion channels, *e.g.*, rod photoreceptor cell channels and olfactory neuron channels that are permeable to cations upon activation by binding of

25 cAMP or cGMP (*see, e.g.*, Altenhofen *et al.*, *PNAS*, 88:9868-72 (1991); Dhallan *et al.*, *Nature*, 347:184-87 (1990)). In cases where activation of the receptor results in a decrease in cyclic nucleotide levels, it may be preferable to expose the cells to agents that increase intracellular cyclic nucleotide levels, *e.g.*, forskolin, prior to adding a receptor-activating compound to the cells in the assay. Cells for this type of assay can

30 be made by co-transfection of a host cell with DNA encoding a cyclic nucleotide-gated ion channel, GPCR phosphatase and DNA encoding a receptor (*e.g.*, certain glutamate receptors, muscarinic acetylcholine receptors, dopamine receptors,

serotonin receptors, and the like), which, when activated, causes a change in cyclic nucleotide levels in the cytoplasm.

In a preferred embodiment, T2R polypeptide activity is measured by expressing a T2R gene in a heterologous cell with a promiscuous G protein that links
5 the receptor to a phospholipase C signal transduction pathway (*see* Offermanns & Simon, *J. Biol. Chem.*, 270:15175-80 (1995)). Optionally the cell line is HEK-293 (which does not naturally express T2R genes) and the promiscuous G protein is G α 15 (Offermanns & Simon, *supra*). Modulation of taste transduction may be assayed by measuring changes in intracellular Ca²⁺ levels, which change in response to
10 modulation of the T2R signal transduction pathway via administration of a molecule that associates with a T2R polypeptide. Changes in Ca²⁺ levels are optionally measured using fluorescent Ca²⁺ indicator dyes and fluorometric imaging.

In one embodiment, the changes in intracellular cAMP or cGMP can be measured using immunoassays. The method described in Offermanns & Simon, *J.*
15 *Bio. Chem.*, 270:15175-180 (1995), may be used to determine the level of cAMP. Also, the method described in Felley-Bosco *et al.*, *Am. J. Resp. Cell and Mol. Biol.*, 11:159-64 (1994), may be used to determine the level of cGMP. Further, an assay kit for measuring cAMP and/or cGMP is described in U.S. Patent 4,115,538, herein incorporated by reference.

20 In another embodiment, phosphatidyl inositol (PI) hydrolysis can be analyzed according to U.S. Patent 5,436,128, herein incorporated by reference. Briefly, the assay involves labeling of cells with ³H-myoinositol for 48 or more hrs. The labeled cells are treated with a test compound for one hour. The treated cells are lysed and extracted in chloroform-methanol-water, after which the inositol phosphates are
25 separated by ion exchange chromatography and quantified by scintillation counting. Fold stimulation is determined by calculating the ratio of cpm in the presence of agonist, to cpm in the presence of buffer control. Likewise, fold inhibition is determined by calculating the ratio of cpm in the presence of antagonist, to cpm in the presence of buffer control (which may or may not contain an agonist).

30 In another embodiment, transcription levels can be measured to assess the effects of a test compound on signal transduction. A host cell containing a T2R polypeptide may be contacted with a test compound for a sufficient time to effect any

interactions, and then the level of gene expression of a protein of interest is measured. The amount of time to effect such interactions may be empirically determined, such as by running a time course and measuring the level of transcription as a function of time. The amount of transcription may be measured by using any method known to those of skill in the art to be suitable. For example, mRNA expression of the protein of interest may be detected using northern blots or their polypeptide products may be identified using immunoassays. Alternatively, transcription based assays using reporter gene may be used as described in U.S. Patent 5,436,128, herein incorporated by reference. The reporter genes can be, *e.g.*, chloramphenicol acetyltransferase, luciferase, '3-galactosidase and alkaline phosphatase. Furthermore, the protein of interest can be used as an indirect reporter via attachment to a second reporter such as green fluorescent protein (*see, e.g.*, Mistili & Spector, *Nature Biotechnology*, 15:961-64 (1997)).

The amount of transcription is then compared to the amount of transcription in either the same cell in the absence of the test compound, or it may be compared with the amount of transcription in a substantially identical cell that lacks the T2R polypeptide. A substantially identical cell may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA. Any difference in the amount of transcription indicates that the test compound has in some manner altered the activity of the T2R polypeptide of interest.

6. Transgenic non-human animals expressing taste receptors

Non-human animals expressing one or more T2R polypeptides of the invention, can also be used for assays. Such expression can be used to determine whether a test compound specifically binds to a mammalian T2R polypeptide *in vivo* by contacting a non-human animal stably or transiently transfected with a nucleic acid encoding a T2R polypeptide or ligand-binding region thereof with a test compound, and determining whether the animal reacts to the test compound by specifically binding to the polypeptide.

Animals transfected or infected with the vectors of the invention are particularly useful for assays to identify and characterize tastants/ligands that can bind to a specific or sets of receptors. Such vector-infected animals expressing human

chemosensory receptor sequences can be used for *in vivo* screening of tastants and their effect on, *e.g.*, cell physiology (*e.g.*, on taste neurons), on the CNS, or behavior.

Means to infect/express the nucleic acids and vectors, either individually or as libraries, are well known in the art. A variety of individual cell, organ, or whole
5 animal parameters can be measured by a variety of means. The T2R sequences of the invention can be for example expressed in animal taste tissues by delivery with an infecting agent, *e.g.*, adenovirus expression vector.

The endogenous chemosensory receptor genes can remain functional and wild-type (native) activity can still be present. In other situations, where it is desirable that
10 all chemosensory receptor activity is by the introduced exogenous hybrid receptor, use of a knockout line is preferred. Methods for the construction of non-human transgenic animals, particularly transgenic mice, and the selection and preparation of recombinant constructs for generating transformed cells are well known in the art.

Construction of a "knockout" cell and animal is based on the premise that the
15 level of expression of a particular gene in a mammalian cell can be decreased or completely abrogated by introducing into the genome a new DNA sequence that serves to interrupt some portion of the DNA sequence of the gene to be suppressed. Also, "gene trap insertion" can be used to disrupt a host gene, and mouse embryonic stem (ES) cells can be used to produce knockout transgenic animals (*see, e.g.*,
20 Holzschu, *Transgenic Res*, 6:97-106 (1997)). The insertion of the exogenous is typically by homologous recombination between complementary nucleic acid sequences. The exogenous sequence is some portion of the target gene to be modified, such as exonic, intronic or transcriptional regulatory sequences, or any genomic sequence which is able to affect the level of the target gene's expression; or a
25 combination thereof. Gene targeting via homologous recombination in pluripotent embryonic stem cells allows one to modify precisely the genomic sequence of interest. Any technique can be used to create, screen for, propagate, a knockout animal, *e.g.*,
see Bijvoet, *Hum. Mol. Genet.*, 7:53-62 (1998); Moreadith, *J. Mol. Med.*, 75:208-16 (1997); Tojo, *Cytotechnology* 19:161-165 (1995); Mudgett, *Methods Mol. Biol.*
30 48:167-184 (1995); Longo, *Transgenic Res*. 6:321-328 (1997); U.S. Patents Nos. 5,616,491; 5,464,764; 5,631,153; 5,487,992; 5,627,059; 5,272,071; WO 91/09955; WO93/09222; WO 96/29411; WO 95/31560; WO 91/12650.

The nucleic acids of the invention can also be used as reagents to produce “knockout” human cells and their progeny. Likewise, the nucleic acids of the invention can also be used as reagents to produce “knock-ins” in mice. The human or rat T2R gene sequences can replace the orthologous T2R in the mouse genome. In this way, a mouse expressing a human or rat T2R is produced. This mouse can then be used to analyze the function of human or rat T2Rs, and to identify ligands for such T2Rs.

F. Modulators

The compounds tested as modulators of a T2R family member can be any small chemical compound, or a biological entity, such as a protein, sugar, nucleic acid or lipid. Alternatively, modulators can be genetically altered versions of a T2R family member. Typically, test compounds may be small chemical molecules and peptides. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays may be designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (*e.g.*, in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, MO), Aldrich (St. Louis, MO), Sigma-Aldrich (St. Louis, MO), Fluka Chemika-Biochemika Analytika (Buchs, Switzerland) and the like.

In one embodiment, high throughput screening methods involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such “combinatorial chemical libraries” or “ligand libraries” are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional “lead compounds” or can themselves be used as potential or actual consumer products.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical “building blocks” such as reagents. For example, a

linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (*i.e.*, the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (*see, e.g.*, U.S. Patent 5,010,175, Furka, *Int. J. Pept. Prot. Res.*, 37:487-93 (1991) and Houghton *et al.*, *Nature*, 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (*e.g.*, WO 91/19735), encoded peptides (*e.g.*, WO 93/20242), random bio-oligomers (*e.g.*, WO 92/00091), benzodiazepines (*e.g.*, U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs *et al.*, *PNAS.*, 90:6909-13 (1993)), vinylogous polypeptides (Hagihara *et al.*, *J. Amer. Chem. Soc.*, 114:6568 (1992)), nonpeptidal peptidomimetics with glucose scaffolding (Hirschmann *et al.*, *J. Amer. Chem. Soc.*, 114:9217-18 (1992)), analogous organic syntheses of small compound libraries (Chen *et al.*, *J. Amer. Chem. Soc.*, 116:2661 (1994)), oligocarbamates (Cho *et al.*, *Science*, 261:1303 (1993)), peptidyl phosphonates (Campbell *et al.*, *J. Org. Chem.*, 59:658 (1994)), nucleic acid libraries (Ausubel, Berger, and Sambrook, all *supra*), peptide nucleic acid libraries (U.S. Patent 5,539,083), antibody libraries (Vaughn *et al.*, *Nature Biotechnology*, 14(3):309-14 (1996) and PCT/US96/10287), carbohydrate libraries (Liang *et al.*, *Science*, 274:1520-22 (1996) and U.S. Patent 5,593,853), small organic molecule libraries (benzodiazepines, Baum, *C&EN*, Jan 18, page 33 (1993); thiazolidinones and metathiazanones, U.S. Patent 5,549,974; pynrolidines, U.S. Patents 5,525,735 and 5,519,134; morpholino compounds, U.S. Patent 5,506,337; benzodiazepines, 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (*see, e.g.*, 357 MPS, 390 MPS (Advanced Chem Tech, Louisville KY), Symphony (Rainin, Woburn, MA), 433A (Applied Biosystems, Foster City, CA), 9050 Plus (Millipore, Bedford, MA)). In addition, numerous combinatorial libraries are themselves commercially available (*see, e.g.*, ComGenex, Princeton, NJ; Tripox,

Inc., St. Louis, MO; 3D Pharmaceuticals, Exton, PA; Martek Biosciences; Columbia, MD; *etc.*).

In one aspect of the invention, the T2R modulators can be used in any food product, confectionery, pharmaceutical composition, or ingredient thereof to thereby modulate the taste of the product, composition, or ingredient in a desired manner. For instance, T2R modulators that enhance bitter taste sensation can be added to provide a bitter taste to a product or composition, while T2R modulators which block bitter taste sensations can be added to improve the taste of a product or composition.

10 **G. Methods for Representing and Predicting the Perception of Taste**

The invention also preferably provides methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the receptors and genes encoding said T2R proteins disclosed herein.

15 Also contemplated as within the invention, is a method of screening one or more compounds for the presence of a taste detectable by a mammal, comprising: contacting said one or more compounds with the disclosed receptors, preferably wherein the mammal is a human. Also contemplated as within the invention, is a method for representing taste perception of a particular taste in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n taste receptors of said vertebrate, where n is greater than or equal to 4; and generating from said values a quantitative representation of taste perception. The taste receptors may be a taste receptor disclosed herein, the representation may constitute a point or a volume in n -dimensional space, may constitute a graph or a spectrum, and may constitute a matrix of quantitative representations. Also, the providing step may comprise contacting a plurality of recombinantly-produced taste receptors with a test composition and quantitatively measuring the interaction of said composition with said receptors.

25 Also contemplated as within the invention, is a method for predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n

taste receptors of said vertebrate, where n is greater than or equal to 4, for one or more molecules or combinations of molecules yielding known taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal, providing values X_1 to X_n representative of the quantitative stimulation of each of n taste receptors of said vertebrate, where n is greater than or equal to 4, for one or more molecules or combinations of molecules yielding unknown taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, and predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal by comparing the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal to the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal. The taste receptors used in this method may include a taste receptor disclosed herein.

In another embodiment, novel molecules or combinations of molecules are generated which elicit a predetermined taste perception in a mammal by determining a value of taste perception in a mammal for a known molecule or combinations of molecules as described above; determining a value of taste perception in a mammal for one or more unknown molecules or combinations of molecules as described above; comparing the value of taste perception in a mammal for one or more unknown compositions to the value of taste perception in a mammal for one or more known compositions; selecting a molecule or combination of molecules that elicits a predetermined taste perception in a mammal; and combining two or more unknown molecules or combinations of molecules to form a molecule or combination of molecules that elicits a predetermined taste perception in a mammal. The combining step yields a single molecule or a combination of molecules that elicits a predetermined taste perception in a mammal.

In another embodiment of the invention, there is provided a method for simulating a taste, comprising the steps of: for each of a plurality of cloned taste receptors, preferably human receptors, ascertaining the extent to which the receptor interacts with the tastant; and combining a plurality of compounds, each having a previously-ascertained interaction with one or more of the receptors, in amounts that together provide a receptor-stimulation profile that mimics the profile for the tastant. Interaction of a tastant with a taste receptor can be determined using any of the binding or reporter assays described herein. The plurality of compounds may then be combined to form a mixture. If desired, one or more of the plurality of the compounds can be combined covalently. The combined compounds substantially stimulate at least 75%, 80%, or 90% of the receptors that are substantially stimulated by the tastant.

In another preferred embodiment of the invention, a plurality of standard compounds are tested against a plurality of taste receptors to ascertain the extent to which the receptors each interact with each standard compound, thereby generating a receptor stimulation profile for each standard compound. These receptor stimulation profiles may then be stored in a relational database on a data storage medium. The method may further comprise providing a desired receptor-stimulation profile for a taste; comparing the desired receptor stimulation profile to the relational database; and ascertaining one or more combinations of standard compounds that most closely match the desired receptor-stimulation profile. The method may further comprise combining standard compounds in one or more of the ascertained combinations to simulate the taste.

H. Kits

T2R genes and their homologs are useful tools for identifying taste receptor cells, for forensics and paternity determinations, and for examining taste transduction. T2R family member-specific reagents that specifically hybridize to T2R nucleic acids, such as T2R probes and primers, and T2R specific reagents that specifically bind to a T2R protein, *e.g.*, T2R antibodies are used to examine taste cell expression and taste transduction regulation.

Nucleic acid assays for the presence of DNA and RNA for a T2R family member in a sample include numerous techniques are known to those skilled in the

art, such as southern analysis, northern analysis, dot blots, RNase protection, S1 analysis, amplification techniques such as PCR, and *in situ* hybridization. In *in situ* hybridization, for example, the target nucleic acid is liberated from its cellular surroundings in such as to be available for hybridization within the cell while
 5 preserving the cellular morphology for subsequent interpretation and analysis. The following articles provide an overview of the art of *in situ* hybridization: Singer *et al.*, *Biotechniques*, 4:230250 (1986); Haase *et al.*, *Methods in Virology*, vol. VII, 189-226 (1984); and Names *et al.*, eds., *Nucleic Acid Hybridization: A Practical Approach* (1987). In addition, a T2R protein can be detected with the various immunoassay
 10 techniques described above. The test sample is typically compared to both a positive control (*e.g.*, a sample expressing a recombinant T2R protein) and a negative control.

The present invention also provides for kits for screening for modulators of T2R family members. Such kits can be prepared from readily available materials and reagents. For example, such kits can comprise any one or more of the following
 15 materials: T2R nucleic acids or proteins, reaction tubes, and instructions for testing T2R activity. Optionally, the kit contains a functional T2R polypeptide. A wide variety of kits and components can be prepared according to the present invention, depending upon the intended user of the kit and the particular needs of the user.

20

EXAMPLES

The following examples provide a summary of the isolated nucleic acid molecules of the invention, and polypeptide sequences corresponding to the conceptual translations of nucleic acid molecules. In the protein sequences presented herein, the one-letter code X or Xaa refers to any of the twenty common amino acid
 25 residues. In the DNA sequences presented herein, the one letter codes N or n refers to any of the of the four common nucleotide bases, A, T, C, or G.

hT2R51 Full-Length cDNA (BAC AC011654) (SEQ ID NO: 1)

30 ATGTTGACTCTAACTCGCATCCGCACTGTGTCCTATGAAGTCAGGAGTACATTTCTGTTCA
 TTTCAGTCCTGGAGTTTGCAGTGGGGTTTCTGACCAATGCCTTCGTTTTCTTGGTGAATTTT
 TGGGATGTAGTGAAGAGGCAGGCACTGAGCAACAGTGATTGTGTGCTGCTGTGCTCAGC
 ATCAGCCGGCTTTTCCTGCATGGACTGCTGTTCTGAGTGCTATCCAGCTTACCCACTTCCA
 35 GAAGTTGAGTGAACCACTGAACCACAGCTACCAAGCCATCATCATGCTATGGATGATTGCA
 AACCAAGCCAACCTCTGGCTTGCTGCCTGCCTCAGCTGCTTTACTGCTCCAAGCTCATCC
 GTTCTCTCACACCTTCTGATCTGCTTGGCAAGCTGGGTCTCCAGGAAGATCTCCAGAT

GCTCCTGGGTATTATTCTTTGCTCCTGCATCTGCACTGTCCTCTGTGTTTGGTGCTTTTTTA
 GCAGACCTCACTTCACAGTCACAACCTGTGCTATTCATGAATAACAATACAAGGCTCAACTG
 GCAGATTAAAGATCTCAATTTATTTTATCTCTCTCTGCTATCTGTGGTCTGTGCCTC
 CTTTCCTATTGTTTCTGGTTTCTTCTGGGATGCTGACTGTCTCCCTGGGAAGGCACATGAGG
 5 ACAATGAAGGTCTATACCAGAAACTCTCGTGACCCAGCCTGGAGGCCACATTAAAGCCC
 TCAAGTCTCTTGTCTCCTTTTTCTGCTTCTTTGTGATATCATCCTGTGTTCCTTCATCTCTG
 TGCCCTACTGATTCTGTGGCGCGACAAAATAGGGGTGATGGTTTGTGTTGGGATAATGGC
 AGCTTGTCCCTCTGGGCATGCAGCCATCCTGATCTCAGGCAATGCCAAGTTGAGGAGAGCT
 GTGATGACCATTCTGCTCTGGGCTCAGAGCAGCCTGAAGGTAAGAGCCGACCACAAGGCA
 10 GATCCCCGACACTGTGCTGA (SEQ ID NO: 1)

hT2R51 Conceptual Translation (BAC AC011654) (SEQ ID NO: 2)

MLTLTRIRTVSYEVRSTFLFISVLEFVAVGFLTNAFVFLVNFWDVVKRQALSNSDCVLLCLSISRL
 15 FLHGLLFLSAIQLTHFQKLSEPLNHSYQAIIMLWMIANQANLWLAACLSLLYCSKLIRFSHTFLI
 CLASWVSRKISQMLLGILCSCICTVLCVWCFFSRPHFTVTTVLFMNNNTRLNWNQKDLNLFYS
 FLFCYLWSVPPFLFLVSSGMLTVSLGRHMRMTMKVYTRNSRDPSEAHKALKSLVSFFCFVVS
 SCVAFISVPLLLWRDKIGVMVCGVIMAAACPSGHAAILISGNAKLRRVMTILLWAQSSLVRA
 20 DHKADSRITC (SEQ ID NO: 2)

hT2R54 Full-Length cDNA (BAC AC024156) (SEQ ID NO: 3)

ATGACTAAACTCTGCGATCCTGCAGAAAGTGAATTGTCGCCATTTCTCATCACCTTAATTTT
 AGCAGTTTTACTTGCTGAATACCTCATTGGTATCATTGCAAATGGTTTCATCATGGCTATAC
 25 ATGCAGCTGAATGGGTTCAAATAAGGCAGTTTCCACAAGTGGCAGGATCCTGGTTTCTCT
 GAGTGTATCCAGAATAGCTCTCCAAAGCCTCATGATGTTAGAAATTACCATCAGCTCAACC
 TCCCTAAGTTTTTATTCTGAAGACGCTGTATATTATGCATTCAAATAAGTTTTATATTCTT
 AAATTTTGTAGCCTGTGGTTTGCTGCCTGGCTCAGTTTCTTCTACTTTGTGAAGATTGCCA
 ATTTCTCCTACCCCTTTTCTCAAAGTGGGAGAAATTACTGGATTGATACCTGGCTT
 30 CTGTGGCTGTCCGTGTTTATTTCTTCAGTCACAGCATGTTCTGCATCAACATCTGCACTGT
 GTATTGTAACAATTTCTTCCCTATCCACTCCTCCAACCTCACTAAGAAAACATACTTGTCTG
 AGATCAATGTGGTGGTCTGGCTTTTCTTTAACTGGGGATTGTGACTCCTCTGATCATG
 TTCATCCTGACAGCCACCCTGCTGATCCTCTCTCTCAAGAGACACACCCTACACATGGGAA
 GCAATGCCACAGGGTCCAACGACCCAGCATGGAGGCTCACATGGGGGCCATCAAAGCTA
 35 TCAGTACTTTCTCATTCTCTACATTTTCAATGCAGTTGCTCTGTTTATCTACCTGTCCAAC
 ATGTTTGACATCAACAGTCTGTGGAATAATTTGTGCCAGATCATCATGGCTGCCTACCCTG
 CCAGCCACTCAATTCTACTGATTCAAGATAACCCTGGGCTGAGAAGAGCCTGGAAGCGGCT
 TCAGCTTCGACTTCATCTTTACCCAAAAGAGTGGACTCTGTGA (SEQ ID NO: 3)

hT2R54 Conceptual Translation (BAC AC024156) (SEQ ID NO: 4)

MTKLCDPAESELSPFLITLILAVLLAEYLIGIANGFIMAIHAAEWVQNKAVSTSGRILVFLSVSRI
 ALQSLMMLEITISSTLSFYSEDAVYYAFKISIFLNFCSLWFAAWLSFFYFVKIANFSYPLFLKL
 45 RWRITGLIPWLLWLSVIFSHSMFCINICTVYCNNSFPIHSSNSTKKTYLSEINVVGLAFFNFLGI
 VTPLIMFILATLLILSLKRHTLHMGSNATGSNDPSMEAHMGAIKAISYFLILYIFNAVALFIYLS
 NMFIDINSLWNNLCQIIMAAYPASHSILLIQDNPGLRRWKRQLRLHLYPKEWTL (SEQ ID
 NO: 4)

hT2R55 Full-Length cDNA (BAC AC024156) (SEQ ID NO: 5)

ATGGCAACGGTGAACACAGATGCCACAGATAAAGACATATCCAAGTTCAAGGTCACCTTC
 ACTTTGGTGGTCTCCGGAATAGAGTGCATCACTGGCATCCTTGGGAGTGGCTTCATCACGG
 CCACTATATGGGGCTGAGTGGGCCAGGGGCAAAACACTCCCCACTGGTGACCGCATATGTT
 GATGCTGAGCTTTTCCAGGCTCTTGCTACAGATTTGGATGATGCTGGAGAACATTTTCAGT
 55 CTGCTATTCCGAATTGTTTATAACCAAAACTCAGTGTATATCCTCTTCAAAGTCATCACTGT
 CTTTCTGAACCATTCGAATCTCTGGTTTGCTGCCTGGCTCAAAGTCTTCTATTGTCTTAGAA
 TTGCAAACTCAATCATCCTTTGTTCTTCTGATGAAGAGGAAAATCATAGTGCTGATGCC

TTGGCTTCTCAGGCTGTCAGTGTGGTTTCCTTAAGCTTCAGCTTTCCTCTCTCGAGAGATG
TCTTCAATGTGTATGTGAATAGCTCCATTCCTATCCCCCTCCTCCAACCTCCACGGAGAAGAA
GTACTTCTCTGAGACCAATATGGTCAACCTGGTATTTTTCTATAACATGGGGATCTTCGTTT
CTCTGATCATGTTTCATCCTGGCAGCCACCCTGCTGATCCTCTCTCTCAAGAGACACACCCTA
5 CACATGGGAAGCAATGCCACAGGGTCCAGGGACCCAGCATGAAGGCTCACATAGGGGCC
ATCAAAGCCACCAGCTACTTCTCATCCTCTACATTTTCAATGCAATTGCTCTATTTCTTTC
CACGTCCAACATCTTTGACACTTACAGTTCCTGGAATATTTTGTGCAAGATCATCATGGCT
GCCTACCCTGCCGGCCACTCAGTACAACTGATCTTGGGCAACCCTGGGCTGAGAAGAGCCT
GGAAGCGGTTTCAGCACCAAGTTCCTCTTTACCTAAAAGGGCAGACTCTGTGA (SEQ ID
10 NO: 5)

hT2R55 Conceptual Translation (BAC AC024156) (SEQ ID NO: 6)

MATVNTDATDKDISKFKVTFTLVVSIGECITGLSGFITAIYGAEWARGKTLPTGDRIMLMLSF
15 SRLLLQIWMMLENIFSLFRIVYNQNSVYILFKVITVFLNHSNLWFAAWLKVFYCLRIANFNHP
LFFLMKRKIIVLPWLLRLSVLSFSFPLSRDVFNVYVNSSIPISSNSTEKKYFSETNMVNLV
FFYNMGIFVPLIMFILAATLLILSLKRHTLHMGSNATGSRDPSMKAHIGAIAKATSYFLILYFNAI
ALFLSTSNIFDTYSSWNILCKIMAAYPAGHSVQLILGNPGLRRWKRQHQVPLYLKGQTL
20 (SEQ ID NO: 6)

hT2R61 Full-Length cDNA (BAC AC018630) (SEQ ID NO: 7)

ATGATAACTTTTCTACCCATCATTTTTCAGTCTGGTAGTGGTTACATTTGTTATTGGAAA
TTTTGCTAATGGCTTCATAGCACTGGTAAATTCATTGAGTGGTTCAAGAGACAAAAGATC
25 TCCTTTGCTGACCAAATTCTCACTGCTCTGGCGGTCTCCAGAGTTGGTTTGGCTCTGGGTATT
ATTATTAACTGGTATTCAACTGTGTTGAATCCAGCTTTTAATAGTGTAGAAGTAAGAACT
ACTGCTTATAATATCTGGGCAGTGATCAACCATTTCAGCAACTGGCTTGCTACTACCCTCA
GCATATTTTATTTGCTCAAGATTGCCAATTTCTCCAACCTTATTTTTCTTCACTTAAAGAGG
AGAGTTAAGAGTGTCAATTCTGGTGATGTTGTTGGGGCCTTTGCTATTTTGGCTTGTCATCT
30 TTTTGTGATAAATGAATGAGATTGTGCGGACAAAAGAATTTGAAGGAAACATGACTTG
GAAGATCAAATTGAAGAGTGCAATGTACTTTTCAAATATGACTGTAACCATGGTAGCAAA
CTTAGTACCCTTCACTCTGACCCTACTATCTTTTATGCTGTTAATCTGTTCTTTGTGTAAAC
ATCTCAAGAAGATGCAGCTCCATGGTAAAGGATCTCAAGATCCCAGCACCAAGGTCCACA
TAAAAGCTTTGCAAACCTGTGATCTCCTTCTCTGTTATGTGCCATTTACTTTCTGTCCATA
35 ATGATATCAGTTTGGAGTTTTGGAAGTCTGGAACAAACCTGTCTTCATGTTCTGCAAAG
CTATTAGATTCACTATCCTTCAATCCACCCATTATCCTGATTTGGGGAAACAAGAAGCT
AAAGCAGACTTTTCTTTCACTTTTGGCAAATGAGGTACTGGGTGAAAGGAGAGAAGACT
TCATCTCCATAG (SEQ ID NO: 7)

40 hT2R61 Conceptual Translation (BAC AC018630) (SEQ ID NO: 8)

MITFLPIIFSSLVVVTFVIGNFANGFIALVNSIEWFKRQKISFADQILTALAVSRVGLLWVLLLNW
YSTVLNPAFNSVEVRTTAYNIWAVINHFSNWLATTLISIFYLLKIANFNSNIFLHLKRRVKSIVLV
MLLGPLLFLACHLFVINMNEIVRTKEFEGNMTWKIKLSAMYFSNMTVTMVANLVPFTLTLLS
45 FMLLICSLCKHLKKMQLHGKGSQDPSTKVHIKALQTVISFLLCAIYFLSIMISVWSFGSLENKP
VFMFCKAIRFSYPSIHPFILIWGNKKLKQTFLSVFWQMRVWVKGEKTSSP (SEQ ID NO: 8)

hT2R63 Full-Length cDNA (BAC AC018630) (SEQ ID NO: 9)

ATGATGAGTTTCTACACATTGTTTTTTCATTCTAGTAGTGGTTGCATTTATTCTTGGA
50 TTTTGCCAATGGCTTTATAGCACTGATAAATTTCAATGCCTGGGTCAAGAGACAAAAGATC
TCCTCAGCTGATCAAATTATTGCTGCTCTGGCAGTCTCCAGAGTTGGTTTGGCTCTGGGTAA
TATTATTACATTGGTATTCAACTGTGTTGAATCCAACCTCATCTAATTTAAAAGTAATAATT
TTTATTTCTAATGCCTGGGCAGTAACCAATCATTTCAAGCATCTGGCTTGCTACTAGCCTCAG
55 CATATTTTATTGCTCAAGATCGTCAATTTCTCCAGACTTATTTTTCATCACTTAAAAAGGA
AGGCTAAGAGTGTAGTTCTGGTGATAGTTGTTGGGGTCTTTGTTCTTTTGGTTTGTACCTT
GTGATGAAACACACGTATATAAATGTGTGGACAGAAGAATGTGAAGGAAACGTAACCTTGG

AAGATCAAACCTGAGGAATGCAATGCACCTTTCCAACCTTGACTGTAGCCATGCTAGCAAACCT
 TGATACCAATTCACCTCTGACCCTGATATCTTTTCTGCTGTTAATCTACTCTCTGTGTAAACAT
 CTGAAGAAGATGCAGCTCCATGGCAAAGGATCTCAAGATCCCAGCACCAAGATCCACATA
 AAAGCTCTGCAAACCTGTGACCTCCTTCCTCATATTACTTGCCATTTACTTTCTGTGTCTAAT
 5 CATATCGTTTTTGAATTTTAAGATGCGACCAAAAAGAAATTGTCTTAATGCTTTGCCAAGCT
 TTTGGAATCATATATCCATCATTCCACTCATTCTGATTTGGGGGAACAAGACGCTAA
 AGCAGACCTTTCTTTCACTTTTGTGGCAGGTGACTTGCTGGGCAAAAGGACAGAACCAGTC
 AACTCCATAG (SEQ ID NO: 9)

10 hT2R63 Conceptual Translation (BAC AC018630) (SEQ ID NO: 10)

MMSFLHIVFSILVVVAFILGNFANGFIALINFIAWVKRQKISSADQIIAALAVSRVGLLWVILLH
 WYSTVLNPTSSNLKVIIFISNAWAVTNHFSIWLATSLSIFYLLKIVNFSRLIFHHLKRKAKSVVLV
 IVLGSIFFLVCHLVMKHTYINVWTECEGNVTWKIKLRNAMHLSNLTVAMLANLIPFTLTLSIF
 15 LLLIYSLCKHLKMKQLHGKGSQDPSTKIHKALQTVTSFLILLAIYFLCLIISFWNFKMRPKEIVL
 MLCQAFGIYPSFHSFILWGNKTLKQTFLSVLWQVTCWAKGQNGSTP (SEQ ID NO: 10)

hT2R64 Full-Length cDNA (BAC AC018630) (SEQ ID NO: 11)

20 ATGACAACTTTTATACCCATCATTTTTTCCAGTGTGGTAGTGGTTCTATTTGTTATTGGAAA
 TTTTGCTAATGGCTTCATAGCATTGGTAAATTCCATTGAGCGGGTCAAGAGACAAAAGATC
 TCTTTTGCTGACCAGATTCTCACTGCTCTGGCGGTCTCCAGAGTTGGTTTGCTCTGGGTATT
 ATTATTAATTTGGTATTCAACTGTGTTAATCCAGCTTTTTATAGTGTAGAAGTAAGAAT
 ACTGCTTATAATGTCTGGGCAGTAACCGGCCATTTCAAGCAACTGGCTTGCTACTGCCCTCA
 25 GCATATTTTATTTGCTCAAGATTGCCAATTTCTCCAACCTTATTTTCTTCACTTAAAGAGG
 AGAGTTAAGAGTGTCTATTCTGGTGATGCTGTTGGGGCCTTTACTATTTTGGCTTGTCAAC
 TTTTGTGTATAAATGAAAGAGATTGTACGGACAAAAGAATATGAAGGAAACTTGACTT
 GGAAGATCAAATTGAGGAGTGCACTGTACCTTTCAAGATGCGACTGTAACCACGCTAGGAA
 ACTTAGTGCCCTTCACTCTGACCCTGCTATGTTTTTGTCTGTTAATCTGTTCTCTGTGTA
 30 CATCTCAAGAAGATGCAGCTCCATGGTAAAGGATCTCAAGATCCCAGCACCAAGGTCCAC
 ATAAAGCTTTGCAAACCTGTGATCTTTTCTCTGTTATGTGCCGTTTACTTTCTGTCCAT
 AATGATATCAGTTTGGAGTTTGGGAGTCTGGAAAACAAACCTGTCTTCATGTTCTGCAAA
 GCTATTAGATTCACTATCCTTCAATCCACCCATTCATCCTGATTTGGGGAAACAAGAAGC
 TAAAGCAGACTTTTCTTTCAGTTTTCGGCAAGTGAGGTACTGGGTGAAAGGAGAGAAGC
 35 CTCATCTCCATAG (SEQ ID NO: 11)

hT2R64 Conceptual Translation (BAC AC018630) (SEQ ID NO: 12)

MTTFPIIFSSVVVVLVIGNFANGFIALVNSIERVKRQKISFADQILTALAVSRVGLLWVLLLNW
 40 YSTVFNPAFYSEVVRTTAYNVWAVTGHFNSWLATSLSIFYLLKIANFSNLIFLHLKRRVKS
 VMLLGPLLFLACQLFVINMKEIVRTKEYEGNLTWKIKLRSAVYLSDATVTTLGNLVPFTLTLLC
 FLLICSLCKHLKMKQLHGKGSQDPSTKVHIKALQTVIFFLLLCVYFLSIMISVWSFGSLENKP
 VFMFCKAIRFSYPSIHPFILWGNKKLKQTFLSVLRQVRYWVKGEKPSSP (SEQ ID NO: 12)

45 hT2R65 Full-Length cDNA (BAC AC018630) (SEQ ID NO: 13)

ATGATGTGTTTTCTGCTCATCATTTTCATCAATTCTGGTAGTGTTCGATTTGTTCTTGGAAA
 TGTTGCCAATGGCTTCATAGCCCTAGTAAATGTCATTGACTGGGTAAACACACGAAAGATC
 TCCTCAGCTGAGCAAATTTCACTGCTCTGGTGGTCTCCAGAATTGGTTTACTCTGGGTCAT
 50 GTTATTCTTTTGGTATGCAACTGTGTTAATTCTGCTTTATATGGTTAGAAAGTAAGAATTG
 TTGCTTCTAATGCCTGGGCTGTAAACGAACCATTTCAAGCATGTGGCTTGCTGCTAGCCTCAG
 CATATTTTGTGCTCAAGATTGCCAATTTCTCCAACCTTATTTCTCTCCACCTAAAGAAGA
 GAATTAAGAGTGTGTTCTGGTGATACTGTTGGGGCCCTTGGTATTTCTGATTTGTAATCTT
 GCTGTGATAACCATGGATGAGAGAGTGTGGACAAAAGAATATGAAGGAAATGTGACTTGG
 55 AAGATCAAATTGAGGAATGCAATACACCTTTCAAGCTTGACTGTAACCTACTCTAGCAAACC
 TCATACCTTTTACTCTGAGCCTAATATGTTTTCTGCTGTTAATCTGTTCTCTTTGTAAACAT
 CTCAAGAAGATGCGGCTCCATAGCAAAGGATCTCAAGATCCCAGCACCAAGGTCCATATA

AAAGCTTTGCAAACCTGTGACCTCCTTCCTCATGTTATTTGCCATTTACTTTCTGTGTATAAT
CACATCAACTTGGAATCTTAGGACACAGCAGAGCAAACCTTGACTCCTGCTTTGCCAAACT
GTTGCAATCATGTATCCTTCATTCCACTCATTATCCTGATTATGGGAAGTAGGAAGCTAA
AACAGACCTTTCTTTCAGTTTGTGGCAGATGACACGCTGA (SEQ ID NO: 13)

5

hT2R65 Conceptual Translation (BAC AC018630) (SEQ ID NO: 14)

MMCFLLISSILVVFVFLGNVANGFIALVNVIDWVNTRKISSAEQILTALVVSRIGLLWVMLFL
WYATVFNSALYGLEVRIVASNAWAVTNHFSMWLAASLSIFCLLKIANFNSNLISHLKKRIKSVV
10 LVILLGPLVFLICNLAVITMDERVWTKEYEGNVTWKIKLRNAIHLSSLTVTTLANLIPFTLSLICF
LLLICSLCKHLKKMRLHSKGSQDPSTKVHIKALQTVTSFLMLFAIYFLCHTSTWNLRTQQSKLV
LLLCQTVAIMYPSFHSFILMGSRKLRKQTFLSVLWQMTR (SEQ ID NO: 14)

hT2R67 Full-Length cDNA (BAC AC018630) (SEQ ID NO: 15)

15

ATGATAACTTTTCTATACATTTTTTTTTTCAATTCTAATAATGGTTTTATTTGTTCTCGGAAA
CTTTGCCAATGGCTTCATAGCACTGGTAAATTTCAATTGACTGGGTGAAGAGAAAAAAGATC
TCCTCAGCTGACCAAATTCCTACTGCTCTGGCGGTCTCCAGAATTGGTTTGCTCTGGGCATT
ATTATTAATTTGGTATTTAACTGTGTTGAATCCAGCTTTTTATAGTGTAGAATTAAGAATT
20 ACTTCTTATAATGCCTGGGTGTAACCAACCATTTCAGCATGTGGCTTGCTGCTAACCTCA
GCATATTTTATTTGCTCAAGATTGCCAATTTCTCCAACCTTCTTTTCTTCATTTAAAGAGG
AGAGTTAGGAGTGTCAATCTGGTGATACTGTTGGGGACTTTGATATTTTGGTTTGTGATC
TTCTTGTTGGCAAACATGGATGAGAGTATGTGGGCAGAAGAATATGAAGGAAACATGACTG
GGAAGATGAAATTGAGGAATACAGTACATCTTTCATATTTGACTGTAACCTACCCTATGGAG
25 CTTCATACCCTTTACTCTGTCCCTGATATCTTTCTGATGCTAATCTGTTCTCTGTGTAAAC
ATCTCAAGAAGATGCAGCTCCATGGAGAAGGATCGCAAGATCTCAGCACCAAGGTCCACA
TAAAAGCTTTGCAAACCTCTGATCTCCTTCTCTGTTATGTGCCATTTTCTTCTATTCTTA
ATCGTTTCGGTTTGGAGTCTAGGAGGCTGCGGAATGACCCGGTTGTCATGGTTAGCAAGG
CTGTTGGAAACATATATCTTGCAATTCGACTCATTATCCTAATTTGGAGAACCAAGAAGCT
30 AAAACACACCTTTCTTTTGATTTGTGTGTCAGATTAGGTGCTGA (SEQ ID NO: 15)

hT2R67 Conceptual Translation (BAC AC018630) (SEQ ID NO: 16)

MITFLYIFFSILIMVFLVFLGNFANGFIALVNFIDWVKRKKISSADQILTALAVSRIGLLWALLLNW
35 YLTVLNPAFYVELRITSYNAWVVTNHFSMWLAANLSIFYLLKIANFNSNLLFLHLKRRVRSVIL
VILLGTLIFLVCHLLVANMDESMWAEYEGNMTGKMKLRNTVHLSYLTVTTLWSFIPFTLSLIS
FLMLICSLCKHLKKMQLHGEQSQDLSTKVHIKALQTLISFLLLCAIFFLFLIVSVWSPRRLRNDP
VVMVSKAVGNIYLAFFDSFILIWRTKKLKHFTLLILCQIRC (SEQ ID NO: 16)

hT2R71 Full-Length cDNA (BAC AC073264) (SEQ ID NO: 17)

ATGCAAGCAGCACTGACGGCCTTCTTCGTGTTGCTCTTTAGCCTGCTGAGTCTTCTGGGGA
TTGCAGCGAATGGCTTCATTGTGCTGGTGCTGGGCAGGGAGTGGCTGCGATATGGCAGGT
TGCTGCCCTTGATATGATCCTCATTAGCTTGGGTGCCTCCCGCTTCTGCCTGCAGTTGGTT
45 GGGACGGTGCACAACTTCTACTACTCTGCCCAGAAGGTCGAGTACTCTGGGGGTCTCGGCC
GACAGTCTTCCATCTACACTGGCACTTCTGAACTCAGCCACCTTCTGGTTTTGCAGCTGG
CTCAGTGTCTGTTCTGTGTGAAGATTGCTAACATCACACACTCCACCTTCTGTGGCTGA
AGTGGAGGTTCCCAGGGTGGGTGCCCTGGCTCCTGTTGGGCTCTGTCCTGATCTCCTTCAT
CATAACCTGCTGTTTTTTTGGGTGAACCTACCCTGTATATCAAGAATTTTAAATTAGAAAAAT
50 TTTCTGGGAACATGACCTACAAGTGGAATACAAGGATAGAAACATACTATTTCCCATCCCT
GAAACTGGTCATCTGGTCAATTCCTTTTTCTGTTTTTCTGGTCTCAATTATGCTGTTAATTA
ATTCTCTGAGGAGGCATACTCAGAGAATGCAGCACACGGGCACAGCCTGCAGGACCCCA
GCACCCAGGCTCACACCAGAGCTCTGAAGTCCCTCATCTCCTTCTCATTCTTTATGCTCTG
TCCTTTCTGTCCCTGATCATTGATGCCGCAAAATTTATCTCCATGCAGAACGACTTTTACTG
55 GCCATGGCAAATTGCAGTCTACCTGTGCATATCTGTCCATCCCTTCATCCTCATCTTCAGCA
ACCTCAAGCTTCGAAGCGTGTCTCGCAGCTCCTGTTGTTGGCAAGGGGCTTCTGGGTGGC
CTAG (SEQ ID NO: 17)

hT2R71 Conceptual Translation (BAC AC073264) (SEQ ID NO: 18)

5 MQAALTAFVLLFSLLSLLGIAANGFIVLVLGREWLRYGRLLPLDMILISLGASRFCLQLVGTVH
NFYYSQAQKVEYSGGLGRQFFHLHWHFLNSATFWFCSWLSVLCVKIANITHSTFLWLKWRFP
WVPWLLLGSVLISFIITLLFFWVNYPVYQEFIRKFSGNMTYKWNTRIETYYFPSLKLVIWSIPFS
VFLVSIMLLINSLRRHTQRMQHNHSLQDPSTQAHTRALKSLISFLILYALSFLSLIIDAAKFISM
QNDFYWPWQIAVYLCISVHPFILIFSNNLKLRSVFSQLLLLLARGFWVA (SEQ ID NO: 18)

10 hT2R75 Full-Length cDNA (SEQ ID NO: 19)

ATGATAACTTTTCTGCCCATCATTTTTTCCATTCTAATAGTGGTTACATTTGTGATTGGAAA
TTTTGCTAATGGCTTCATAGCATTGGTAAATTCATTGAGTGGTTCAAGAGACAAAAGATC
TCTTTTGCTGACCAAATCTCACTGCTCTGGCAGTCTCCAGAGTTGGTTTACTCTGGGTATT
15 AGTATTAAATGGTATGCAACTGAGTTGAATCCAGCTTTTAACAGTATAGAAGTAAGAATT
ACTGCTTACAATGTCTGGGCAGTAATCAACCATTTCAGCAACTGGCTTGCTACTAGCCTCA
GCATATTTTATTTGCTCAAGATTGCCAATTTCTCCAACCTTATTTTTCTTCACTTAAAGAGG
AGAGTTAAGAGTGTTGTTCTGGTGATACTATTGGGGCCTTTGCTATTTTTGGTTTGTCTATCT
TTTTGTGATAAACATGAATCAGATTATATGGACAAAAGAATATGAAGGAAACATGACTTG
20 GAAGATCAAACCTGAGGAGTGCAATGTACCTTTCAAATACAACGGTAACCATCCTAGCAAA
CTTAGTTCCTTCACTCTGACCCTGATATCTTTTCTGCTGTTAATCTGTTCTCTGTGTAAAC
ATCTCAAAAAGATGCAGCTCCATGGCAAAGGATCTCAAGATCCCAGCATGAAGGTCCACA
TAAAGCTTTGCAAACCTGTGACCTCCTTCTCTGTTATGTGCCATTTACTTTCTGTCCATA
ATCATGTCAAGTTTGGAGTTTGGAGAGTCTGGAAAACAAACCTGTCTTCTCATGTTCTGCGAAG
25 CTATTGCATTCACTATCCTTCAACCCACCCATTATCCTGATTTGGGGAAACAAAGAGCT
AAAGCAGACTTTTCTTTCAGTTTGTGGCATGTGAGGTACTGGGTGAAAGGAGAGAAGCCT
TCATCTTCATAG (SEQ ID NO: 19)

hT2R75 Conceptual Translation (SEQ ID NO: 20)

30 MITFLPIIFSLIVVTFVIGNFANGFIALVNSIEWFKRQKISFADQILTALAVSRVGLLWVLVLNW
YATELNPAPNSIEVRITAYNVWAVINHFSNWLATSLSIFYLLKIANFSNLIFLHLKRRVKSIVLVI
LLGPLLVCHLVFINMNQIIWTKKEYEGNMTWKIKLRSA MYLSNNTTVTILANLVPFTLTLISFLL
LICSLCKHLKMKQLHGKGSQDPSMKVHIKALQTVTSFLLCAIYFLSIMSVWSFESLENKPVF
35 MFCBAIAFSYPSTHPFILIWGNKKLKQTFLSVLWHVRYWVKGEKPSSS (SEQ ID NO: 20)

hT2R59 Pseudogene (BAC AC018630) (SEQ ID NO: 21)

ATGGTATATTTTCTGCTCATCATTTTTATCAATTCTGGTAGTGGTTGCATTTGTTCTTGAAAA
40 TTTTTCGAATGGCTTCATAGCTCTAGTAAATGTCATTGACTGGGTAAAGACACGAAAGATC
TCCTCAGCTGACCAAATCCTCACTGCTCTGGTGGTCTCCAGAATTGGTTTACTCTGGGTCTAT
ATTATTACATTGGTATGCAAATGTGTTAATTCAGCTTTATATAGTTCAAGAGTAGGAGCT
GTTGCTTCTAATATCTCAGCAATAATCAACCATTTCAGCATCTGGCTTGCTGCTAGCCTCAG
CATATTTTATTTGCTCAAGATTGCCAATTTCTCCAACCTTATTTTTCTCCACCTAAAGAGA
45 GAATTAGGAGTGTTGTTCTGGTGATACTGTTGGGTCCCTTGGTATTTTTGATTTGTAATCTT
GCTGTGATAACCATGGATGACAGTGTGTGGACAAAAGAATATGAAGGAAATGTGACTTGG
AAGATCAAATTGAGGAATGCAATACACCTTTCAAACCTGACTGTAAGCACACTAGCAAACC
TCATACCTTCACTCTGACCCTAATATGTTTTCTGCTGTTAATCTGTTCTCTGCATAAACAT
CTCAAGAAGATGCAGCTCCATGGCAAAGGATCTCAAGATCTCAGCACCAAGGTCCACATA
50 AAAGCTTTGCAAACCTGTGATCTCCTTCTCATGTTATATGCCATTTACTTTCTGTATCTAAT
CACATTAACCTGGAATCTTTGAACACAGCAGAACAACTTGATTTCTGCTTTGCAAACCT
CTTGGAATCATGTATCCTTCACTTCACTCATTCTTCTGATTATGGGAAGCAGGAAACTAA
AACAGACGTTTCTTTCAGTTTATGTGAGGTACATGCTTAGTGAAAGGACAGCAACCCTC
AACTCCATAG (SEQ ID NO: 21)

55

hT2R69 Pseudogene (BAC AC018630) (SEQ ID NO: 22)

5 ATGATATGTTTTCTGCTCATCATTTTATCAATTCTGGTAGTGTTCATTGTTCTTGGAAA
 TGTGTTGCCAATGGCTTCATAGCTCTAGTAGGTGTCCTTGAGTGGGTAAAGACACAAAAGATC
 TCATCAGCTGACCAAATTTCTCACTGCTCTGGTGGTGTCCAGAGTTGGTTTACTCTGGGTC
 ATATTATTACATTGGTATGCAACTGTGTTAATTTGGCTTCACATAGATTAGAAGTAAGAA
 TTTTGGTTCTAATGTCTCAGCAATAACCAAGCATTTCAGCATCTGGGTGTTACTAGCCTCA
 GCATATTTTCATTTGCTCAAGACTGCCAATTTCTCCAACCTTATTTTCTCCACCTAAAGAAA
 10 AGGATTAAGAATGTTGGTTTGGTGATGCTGTTGGGGCCCTTGGTATTTTTCATTGTAAATC
 TTGCTCTGATAACCAACGGGTGAGAGTGTGTGGACAAAAGAATATGAAGGAAATTTGTCTT
 GGATGATCAAATTGAGGAATGCAATACAGCTTTCAAACCTTGACTGTAACCATGCCAGCAA
 ACGTCACACCCTGCACTCTGACACTAATATCTTTCTGCTGTAAATCTATTCTCCATGTAAA
 CATGTCAAGAAGATGCAGCTCCATGGCAAAGGATCTCAACATCTCAGCACCAAGGTGCAC
 AATAAAGCTTTGCAAACGTGTATCTCCTTCCTTATGTTATTTGCCATTTACTTTCTGTGTCT
 15 AATCACATCAACTTGGAAATCCTAGGACTCAGCAGAGCAAACCTTGATTCTCTGCTTTACCAA
 ACTCTTGGATTCAATGTATCTTTGTTCCACTCATTATCTGACTATGGGAAGTAGGAAGCC
 AAAACAGACCTTTCTTTCAGCTTTGTGA (SEQ ID NO: 22)

mT2R33 Full-Length cDNA (BAC AC020619) (SEQ ID NO: 23)

20 ATGACCTCCCCTTTCCAGCTATTTATCATATGGTCATCATGACAGCAGAGTTTCTCATCGG
 GACTACAGTGAATGGATTCCCTTATCATTGTGAAGTCTATGACTTGTTCAGAGCCGAACG
 TTCTGATCCTGCAGACCCTCTTGATGTGCACAGGGCTGTCCAGACTCGGTCTGCAGATAA
 TGCTCATGACCCAAAGCTTCTTCTGTGTTCTTTCCATACTCTTATGAGGAAAATATTTAT
 25 AGTTCAGATATAATGTTCTGCTGGATGTTCTTCAGCTCGATTGGCCTCTGGTTTGCCACATG
 TCTCTCTGTCTTTACTGCCTCAAGATTTCAAGCTTCACTCCACCCTGGTTTCTTTGGCTGA
 AATTCAGAATTTCAAAGCTCATATTTGGCTGCTTCTGGGCAGCTTGCTGGCCTCTCTGGG
 CACTGCAACTGTGTGCATCGAGGTAGGTTTCCCTTTAATTGAGGATGGCTATGTCCTGAGA
 AACGCAGGACTAAATGATAGTAATGCCAAGCTAGTGAGAAATAATGACTTGCTCCTCATC
 30 AACCTGATCCTCCTGCTTCCCCTGTCTGTGTTTGTGATGTGCACCTCTATGTTATTTGTTTC
 TCTTTACAAGCATGCACTGGATGCAAAGCGAATCTCACAAGCTGTCAAGTGCCAGAACC
 GAAGCTCATATAAATGCATTAAAGACAGTGACAACATTCCTTTGTTTCTTTGTTTCTTACTT
 TGCTGCCTTCATGGCAAATATGACATTTAGAATTCACATACAGAAGTCATCAGTTCCTCGTG
 GTGAAGGAAATCATGGCAGCATATCCCGCCGGCCACTCTGTCTACATAATCGTCTTGAGTAACT
 35 CTAAGTTCAAAGACTTATTCAGGAGAATGATCTGTCTACAGAAGGAAGAGTGA (SEQ ID
 NO: 23)

mT2R33 Conceptual Translation (BAC AC020619) (SEQ ID NO: 24)

40 MTSPPFAIYHVMIMTAEFLLGTTVNGFLIIVNCYDLFKSRTFLILQTLMLMCTGLSRLGLQIMLMT
 QSFFSVFFPYSEENIYSSDIMFVWMFFSSIGLWFATCLSVFYCLKISGFTPPWFLWLKFRISKLI
 WLLGSLLASLGTATVCIEVGFLIEDGYVLRNAGLNDNSNAKLVRNNDLLLNLILLPLSVFVM
 CTSMLFVSLYKMHWMQSESHKLSSARTEAHINALKTVTTFFCFVSFYFAAFMANMTFRIPYR
 45 SHQFFVVKEMAAYPAGHSVIIVLSNSKFKDLFRRMICLQKEE (SEQ ID NO: 24)

While the foregoing detailed description has described several embodiments of
 the present invention, it is to be understood that the above description is illustrative
 only and not limiting of the disclosed invention. The invention is to be limited only
 by the claims which follow.

WHAT IS CLAIMED:

1. An isolated nucleic acid molecule selected from the group consisting of:

(i) an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof which comprises at least 75 nucleotides;

(ii) an isolated cDNA or an insoluble RNA transcribed therefrom that encodes a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which encodes at least 25 contiguous amino acids of said polypeptide;

(iii) an isolated nucleic acid molecule having at least 20-30% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or to a fragment thereof which comprises at least 100 contiguous nucleotides thereof;

(iv) an isolated nucleic acid molecule that encodes a polypeptide having at least 40% sequence identity at the amino acid level with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or an isolated nucleic acid molecule encoding at least 50 contiguous amino acid residues thereof;

(v) an isolated nucleic acid molecule that encodes a taste receptor or a fragment thereof that specifically hybridizes under stringent hybridization conditions to a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23; and

(vi) a variant of an isolated nucleic acid molecule according to (i) or (ii), containing at least one substitution, deletion or addition mutation in the coding region.

2. The isolated nucleic acid molecule of Claim 1 which is selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof which comprises at least 75 contiguous nucleotides thereof.

3. The isolated nucleic acid molecule of Claim 1 which encodes a polypeptide having an amino acid sequence selected from the group consisting of SEQ

ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof encoding at least 25 contiguous amino acid residues of said polypeptide.

4. An isolated nucleic acid molecule having at least 30% sequence
5 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof comprising at least 100 contiguous nucleotides of any of said sequences.

5. An isolated nucleic acid molecule having at least 40-60% sequence
10 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof comprising at least 100 contiguous nucleotides of any of said sequences.

6. An isolated nucleic acid molecule having at least 70% sequence
15 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof comprising at least 100 contiguous nucleotides of any of said sequences.

7. An isolated nucleic acid molecule having at least 80% sequence
20 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof comprising at least 100 contiguous nucleotides of any of said sequences.

8. An isolated nucleic acid molecule having at least 90% sequence
25 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or a fragment thereof comprising at least 100 contiguous nucleotides of any of said sequences.

9. An isolated nucleic acid molecule according to Claim 1 which encodes
30 a polypeptide having at least 95% sequence identity with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8,

10, 12, 14, 16, 18, 20, and 24, or a fragment thereof that comprises at least 40 contiguous amino acids thereof.

10. An isolated nucleic acid molecule according to Claim 1 which encodes
5 a polypeptide having at least 96% sequence identity with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof that comprises at least 40 contiguous amino acids thereof.

10. 11. An isolated nucleic acid molecule according to Claim 1 which encodes a polypeptide having at least 97% sequence identity with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof that comprises at least 40 contiguous amino acids thereof.

15

12. An isolated nucleic acid molecule according to Claim 1 which encodes a polypeptide having at least 98% sequence identity with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof that comprises at least 40
20 contiguous amino acids thereof.

13. An isolated nucleic acid molecule according to Claim 1 which encodes a polypeptide having at least 99% sequence identity with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8,
25 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof that comprises at least 40 contiguous amino acids thereof.

14. An isolated nucleic acid molecule which exhibits at least 20-30% sequence identity with a nucleic acid sequence selected from the group consisting of
30 SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23 or which exhibits at least 30% sequence identity to a fragment comprising at least 100 contiguous nucleotides of said nucleic acid sequence; and which further contains at least one sequence that encodes a

polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

15 15. An isolated nucleic acid molecule which exhibits at least 40-60%
sequence identity with a nucleic acid sequence selected from the group consisting of
SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or which exhibits at least 40-
60% sequence identity to a fragment comprising at least 100 contiguous nucleotides of
said nucleic acid sequence; and which further contains at least one sequence that
encodes a polypeptide which is at least 70% identical to a consensus sequence
10 selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

16. An isolated nucleic acid molecule that exhibits at least 70% sequence
identity with a nucleic acid sequence selected from the group consisting of SEQ ID
NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or at least 70% sequence identity with a
15 fragment thereof comprising at least 100 contiguous nucleotides thereof and; which
further contains at least one sequence that encodes a polypeptide which is at least 70%
identical to a consensus sequence selected from the group consisting of SEQ ID NOS:
25, 26, 27, 28, 29, and 30.

20 17. An isolated nucleic acid molecule that exhibits at least 80% sequence
identity with a nucleic acid sequence selected from the group consisting of SEQ ID
NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or at least 90% sequence identity with a
fragment thereof comprising at least 100 contiguous nucleotides thereof; and which
further contains at least one sequence that encodes a polypeptide which is at least 70%
25 identical to a consensus sequence selected from the group consisting of SEQ ID NOS:
25, 26, 27, 28, 29, and 30.

18. An isolated nucleic acid molecule that exhibits at least 85% sequence
identity with a nucleic acid sequence selected from the group consisting of SEQ ID
30 NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or at least 85% sequence identity with a
fragment thereof comprising at least 100 contiguous nucleotides thereof; and which
further contains at least one sequence that encodes a polypeptide which is at least 70%

identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

19. An isolated nucleic acid molecule that exhibits at least 90% sequence
5 identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or at least 90% sequence identity with a fragment thereof comprising at least 100 contiguous nucleotides thereof; and which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS:
10 25, 26, 27, 28, 29, and 30.

20. An isolated nucleic acid molecule that exhibits at least 95% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or at least 95% sequence identity with a
15 fragment thereof comprising at least 100 contiguous nucleotides thereof; and which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

21. An isolated nucleic acid molecule that exhibits about 95-99% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23, or about 95-99% sequence identity with a fragment thereof comprising at least 100 contiguous nucleotides thereof; and which further contains at least one sequence that encodes a polypeptide which is at least 70%
25 identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

22. An isolated nucleic acid molecule which encodes for a functional taste receptor that comprises a portion which is at least 100 nucleotides in length that
30 exhibits at least 40% sequence identity with at least 100 contiguous nucleotides of a portion of a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23.

23. The nucleic acid molecule of Claim 22 which is a chimeric nucleic acid molecule, wherein said chimeric nucleic acid molecule is produced by combining portions of at least two different G Protein-Coupled Receptors.

5

24. The chimeric nucleic acid molecule of Claim 23, wherein said two different G Protein-Coupled Receptors are taste receptors.

25. The chimeric nucleic acid molecule of Claim 23, wherein said chimeric
10 nucleic acid molecule contains at least 200 contiguous nucleotides that are at least 40% identical to a portion of one of said nucleic acid sequences.

26. An isolated nucleic acid molecule according to Claim 1 which is
15 directly or indirectly attached to a nucleic acid sequence that encodes a detectable polypeptide.

27. The nucleic acid molecule of Claim 26, wherein said detectable polypeptide is green fluorescent protein, or a fragment or variant thereof.

20 28. An isolated nucleic acid molecule which encodes a polypeptide that exhibits about 30-40% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of
25 said polypeptide on the surface of a cell.

29. An isolated nucleic acid molecule which encodes a polypeptide that exhibits about 50% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment
30 thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

30. An isolated nucleic acid molecule which encodes a polypeptide that exhibits about 60% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

31. An isolated nucleic acid molecule which encodes a polypeptide that exhibits at least 70% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

15

32. An isolated nucleic acid molecule which encodes a polypeptide that exhibits at least 80% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

33. An isolated nucleic acid molecule which encodes a polypeptide that exhibits at least 90% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

34. An isolated nucleic acid molecule which encodes a polypeptide that exhibits at least 95% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment

thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of said polypeptide on the surface of a cell.

5 35. An isolated nucleic acid molecule which encodes a polypeptide that exhibits about 95 to 99% sequence identity with a polypeptide selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof comprising at least 40 contiguous amino acids thereof that is directly or indirectly attached to a sequence that facilitates the expression and/or translocation of
10 said polypeptide on the surface of a cell.

 36. An isolated nucleic acid molecule according to Claim 1 which is operably linked to a constitutive promoter.

15 37. An isolated nucleic acid molecule according to Claim 1 which is operably linked to a regulatable promoter.

 38. An isolated nucleic acid molecule according to Claim 1 which is directly or indirectly attached to a nucleic acid sequence encoding a chaperone protein
20 or a fragment thereof.

 39. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a fragment of at least 60 contiguous amino acids of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8,
25 10, 12, 14, 16, 18, 20, and 24.

 40. The isolated nucleic acid molecule of Claim 39, wherein the nucleotide sequence encodes at least 100 amino acids.

30 41. The isolated nucleic acid molecule of Claim 39, wherein the nucleotide sequence encodes at least 150 amino acids.

42. The isolated nucleic acid molecule of Claim 39 wherein the nucleotide sequence encodes at least 200 amino acids.

43. The isolated nucleic acid molecule of Claim 39, wherein the nucleotide
5 sequence encodes at least 250 amino acids.

44. The isolated nucleic acid molecule of Claim 39, wherein the polypeptide is a T2R polypeptide

10 45. The isolated nucleic acid molecule of Claim 39, wherein the fragment specifically binds a ligand selected from the group consisting of 6-n-propylthiouracil, sucrose octaacetate, raffinose undecaacetate, cycloheximide, denatonium, copper glycinate, and quinine.

15 46. An isolated nucleic acid molecule according to Claim 1 comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, and 23.

47. An expression vector that comprises a nucleic acid sequence according
20 to Claim 1.

48. The expression vector of Claim 47 which is a mammalian, yeast, bacterial or insect expression vector.

25 49. A cell which is transfected or transformed with at least one nucleic acid sequence according to Claim 1.

50. A cell according to Claim 49, wherein the cell is mammalian.

30 51. A mammalian cell according to Claim 50, wherein the cell human.

52. A cell according to Claim 49, wherein the cell is yeast or insect.

53. The mammalian cell according to Claim 50 which is selected from the group consisting of a taste cell, Chinese hamster ovary cell, baby hamster kidney cell, and a myeloma cell.

5

54. A solid phase comprising at least one isolated nucleic acid sequence according to Claim 1.

55. A solid phase which is attached to an array of different nucleic acid sequences which includes at least one sequence according to Claim 1.

10

56. The solid phase according to Claim 54 which comprises an array of at least 4 different nucleic acid sequences that encode functional taste receptors or fragments or variants thereof.

15

57. A solid phase comprising the nucleic acid array of Claim 54 which comprises at least 10 different nucleic acid sequences that encode functional taste receptors.

58. A solid phase comprising the nucleic acid array of Claim 54 which comprises at least 50 different nucleic acid sequences that encode taste receptors or fragments or variants thereof.

20

59. A solid phase comprising the nucleic acid array of Claim 54 which comprises at least 100 different sequences that encode taste receptors or fragments or variants thereof.

25

60. An isolated polypeptide or fusion protein that is selected from the group consisting of:

(i) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24;

30

(ii) a polypeptide comprising an amino acid sequence that exhibits at least 50% sequence identity with an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24;

(iii) a polypeptide comprising an amino acid sequence that exhibits at least 75% sequence identity with a fragment of a polypeptide according to (i) that is at least 40 amino acids in length;

(iv) a chimeric polypeptide that comprises a portion of a polypeptide according to (i) or (ii) that is at least 40 amino acids in length, and a portion of at least one other G Protein-Coupled Receptor; and

(v) a variant of a polypeptide according to (i) which differs by said polypeptide by at least one substitution, addition or deletion modification.

61. An isolated polypeptide according to Claim 60 wherein such polypeptide exhibits at least 60% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

62. An isolated polypeptide according to Claim 60 wherein said polypeptide exhibits at least 65% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

63. An isolated polypeptide according to Claim 60 wherein said polypeptide exhibits at least 75% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

64. An isolated polypeptide according to Claim 60 wherein said polypeptide exhibits at least 75% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

65. An isolated polypeptide according to Claim 60 wherein said polypeptide exhibits at least 80% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

5

66. An isolated polypeptide according to Claim 60 wherein said polypeptide exhibits at least 90% sequence identity with a polypeptide having a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24, or a fragment thereof which is at least 200 amino acids.

10

67. A variant according to Claim 60(v) which comprises at least 5 conservative amino acid substitutions.

68. A variant according to Claim 60(v) which comprises at most 5 conservative amino acid substitutions.

15

69. A variant according to Claim 60(v) which comprises 5 to 7 conservative substitution modifications.

70. A variant according to Claim 60(v) which comprises 3 to 4 conservative substitution modifications.

20

71. A variant according to Claim 60(v) which comprises 1 or 2 conservative substitution modifications.

25

72. A solid phase to which is immobilized directly or indirectly at least one isolate polypeptide according to Claim 60 or a cell which expresses said polypeptide on the surface thereof.

73. The solid phase of Claim 72 which comprises immobilized at least 4 different polypeptides according to Claim 60 or a cell which expresses said polypeptide on the surface thereof.

30

74. The solid phase of Claim 72 which comprises immobilized at least 16 different polypeptides according to Claim 60 or a cell which expresses said polypeptide on the surface thereof.

5

75. The solid phase of Claim 72 which comprises immobilized at least 25 different polypeptides according to Claim 48 or a cell which expresses said polypeptide on the surface thereof.

10 76. A method of detecting expression of a taste receptor gene comprising (a) hybridizing at least one sample with a nucleic acid molecule according to Claim 1, and (b) detecting expression of the taste receptor gene by a positive hybridization signal.

15 77. A method of screening a library comprising (a) hybridizing the library with a nucleic acid molecule according Claim 1 and (b) detecting one or more taste receptor clones in the library by a positive hybridization signal.

20 78. A recombinant polynucleotide comprising a nucleic acid molecule according to Claim 1 which is attached directly or indirectly to a heterologous nucleic acid molecule.

25 79. An expression vector comprising a nucleic acid molecule according to Claim 1, wherein the nucleic acid molecule is operably linked to a heterologous nucleic acid molecule that drives expression thereof.

80. A transfected or transformed cell comprising the recombinant polynucleotide of Claim 78 introduced into a host cell, or a progeny thereof.

30 81. A transgenic non-human organism comprising the recombinant polynucleotide of Claim 78 introduced into a cell of a host non-human organism, or a progeny thereof.

82. A method of making a recombinant polynucleotide comprising ligating a nucleic acid molecule according to Claim 1 to a heterologous nucleic acid.

5 83. The method of Claim 82 wherein the heterologous nucleic acid comprises a translational and/or transcriptional regulatory region.

84. A method of making a transfected cell comprising introducing the recombinant polynucleotide of Claim 78 into a host cell, or propagating the host cell
10 in which the recombinant polynucleotide has been introduced.

85. A method of detecting specific binding of a putative ligand to a taste receptor comprising (a) contacting the putative ligand with a cell in which the expression vector of Claim 79 has been introduced, wherein the taste receptor is
15 expressed by the cell thereby, and (b) directly or indirectly detecting specific binding between the putative ligand and the taste receptor.

86. A method of making transgenic non-human organism comprising introducing the recombinant polynucleotide of Claim 78 into a cell of a host non-
20 human organism, or propagating the host non-human organism in which the recombinant polynucleotide has been introduced.

87. An isolated polypeptide molecule comprising a fragment of at least 60 contiguous amino acids of a polypeptide having an amino acid sequence selected from
25 the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24.

88. The isolated polypeptide molecule of Claim 87, wherein the fragment contains at least 100 amino acids.

30 89. The isolated polypeptide molecule of Claim 87, wherein the fragment contains at least 150 amino acids.

90. The isolated polypeptide molecule of Claim 87, wherein the fragment contains at least 200 amino acids.
91. The isolated polypeptide molecule of Claim 87, wherein the fragment
5 contains at least 250 amino acids.
92. The isolated polypeptide molecule of Claim 87, wherein the fragment is a T2R polypeptide.
93. The isolated polypeptide molecule of Claim 87, wherein the fragment
10 specifically binds a tastant associated with a bitter substance selected from the group consisting of 6-n-propylthiouracil, sucrose octaacetate, raffinose undecaacetate, cycloheximide, denatonium, copper glycinate, and quinine.
94. A recombinant polypeptide comprising the polypeptide of Claim 87
15 and a heterologous peptide domain.
95. The recombinant polypeptide of Claim 94, wherein the heterologous peptide domain comprises a transmembrane region of a G Protein-Coupled Receptor.
20
96. The recombinant polypeptide of Claim 94 comprising a seven-transmembrane receptor with a taste receptor ligand-binding region, wherein the taste receptor ligand-binding region is a chimera of at least two different taste receptors.
97. A method of detecting specific binding of a ligand to a taste receptor
25 comprising (a) contacting the ligand with the polypeptide of Claim 87, and (b) directly or indirectly detecting specific binding between the ligand and the taste receptor.
98. An antibody or antibody fragment that specifically binds a polypeptide
30 having an amino acid sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24.

99. A method of detecting specific binding of the antibody of Claim 98 to a taste receptor comprising (a) contacting the antibody with a sample which may contain the taste receptor and (b) detecting specific binding therebetween.

5 100. The method of Claim 99, wherein specific binding of the antibody to a cell in the sample identifies the cell as a taste cell.

101. A method of screening a library of chemical compounds for compounds that are involved in taste sensation comprising contacting compounds in
10 said library with at least one polypeptide according to Claim 87 and identifying compounds that specifically bind to at least one of said polypeptides.

102. The method of Claim 101 wherein said library is a combinatorial chemical library.

15

103. The method of Claim 101 wherein said library is a peptide library.

104. The method of Claim 101 wherein said library is a peptide, encoded peptide, benzodiazepine, diversomer, vinylogous polypeptide, nonpeptidal
20 peptidominetic, or small molecule organic compound library.

105. The method of Claim 101 wherein said library is a random combination of compounds.

25 106. The method of Claim 101 wherein said compounds are screened by high throughput screening.

107. The method of Claim 101 wherein said screening is effected using animal cells or tissues that express at least one of said polypeptides.

30

108. A cell or non-human animal assay for identifying molecules that interact with a T2R polypeptide comprising:

obtaining a cell or non-human animal that expresses at least one functional T2R polypeptide or a chimeric polypeptide comprising a functional T2R polypeptide, and which optionally expresses at least one functional G Protein;

contacting said cell or non-human animal with a molecule to be screened for
5 its ability to modulate a T2R polypeptide; and
detecting whether modulation occurs.

109. The method of Claim 108 wherein modulation is detected based on changes in intracellular calcium.

10

110. The method of Claim 108 wherein modulation is detected by measuring the transfer of ^{32}P from gamma-labeled GTP to the T2R polypeptide.

111. The method of Claim 108 wherein modulation is determined based on
15 a comparison to a control compound known to modulate the particular T2R polypeptide.

112. The method of Claim 108 wherein the G Protein is G α 15 or G α 16 or another promiscuous G Protein.

20

113. The method of Claim 108 wherein modulation is determined by detecting whether a change in the level of intracellular cyclic nucleotides occurs.

114. The method of Claim 108 wherein modulation is determined based on
25 the level of transcription a predetermined target protein after contacting the cell with the screened compound.

115. The method of Claim 108 when said screened compounds are synthesized by computer assisted compound design based on the predicted or actual
30 three-dimensional structure of the amino acid sequence of the T2R polypeptide or a fragment thereof.

116. The method of Claim 108 wherein compounds that modulate T2R polypeptides are identified based on whether they specifically bind to a T2R polypeptide.

5 117. The method of Claim 108 wherein modulation refers to the inhibition of T2R polypeptide function.

118. The method of Claim 108 wherein modulation refers to the enhancement of T2R polypeptide function.

10

119. A method for representing the perception of one or more tastes in one or more mammals, comprising:

providing values X_1 to X_n representative of the quantitative stimulation of each of n taste receptors of said mammals; and

15 generating from said values a quantitative representation of taste perception, wherein at least one of said taste receptors is a taste receptor polypeptide having a sequence that is at least about 75% identical to a sequence selected from the group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24.

20 120. The method of claim 119, wherein said representation constitutes a point or a volume in n -dimensional space.

121. The method of claim 119, wherein said representation constitutes a graph or a spectrum.

25

122. The method of claim 119, wherein said representation constitutes a matrix of quantitative representations.

30 123. The method of claim 119, wherein said providing step comprises contacting a plurality of recombinantly produced taste receptors with a test composition and quantitatively measuring the interaction of said composition with said receptors.

124. A method for predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules comprising:
- providing values X_1 to X_n representative of the quantitative stimulation of each
- 5 of n taste receptors of said mammal, for one or more molecules or combinations of molecules yielding known taste perception in a mammal,
- generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal;
- 10 providing values X_1 to X_n representative of the quantitative stimulation of each of n taste receptors of said mammal, for one or more molecules or combinations of molecules yielding unknown taste perception in a mammal;
- generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding
- 15 unknown taste perception in a mammal; and
- predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal by comparing the quantitative representation of taste perception in a mammal generated by one or more molecules or combinations of molecules yielding
- 20 unknown taste perception in a mammal to the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal,
- wherein at least one of said taste receptors is a taste receptor polypeptide having a sequence that is at least about 75% identical to a sequence selected from the
- 25 group consisting of SEQ ID NOS: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, and 24.

125. An isolated nucleic acid molecule which exhibits at least 20-30% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 21 and 22, or which exhibits at least 30% sequence identity to a
- 30 fragment comprising at least 100 contiguous nucleotides of said nucleic acid sequence; and which further contains at least one sequence that encodes a polypeptide

which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

126. An isolated nucleic acid molecule which exhibits at least 40-60% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 21 and 22, or which exhibits at least 40-60% sequence identity to a fragment comprising at least 100 contiguous nucleotides of said nucleic acid sequence; and which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

127. An isolated nucleic acid molecule that exhibits at least 70% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 21 and 22, or at least 70% sequence identity with a fragment thereof comprising at least 100 contiguous nucleotides thereof and; which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

128. An isolated nucleic acid molecule that exhibits at least 80% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 21 and 22, or at least 90% sequence identity with a fragment thereof comprising at least 100 contiguous nucleotides thereof; and which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

129. An isolated nucleic acid molecule that exhibits at least 85% sequence identity with a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 21 and 22, or at least 85% sequence identity with a fragment thereof comprising at least 100 contiguous nucleotides thereof; and which further contains at least one sequence that encodes a polypeptide which is at least 70% identical to a consensus

sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

130. An isolated nucleic acid molecule that exhibits at least 90% sequence
5 identity with a nucleic acid sequence selected from the group consisting of SEQ ID
NOS: 21 and 22, or at least 90% sequence identity with a fragment thereof comprising
at least 100 contiguous nucleotides thereof; and which further contains at least one
sequence that encodes a polypeptide which is at least 70% identical to a consensus
sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and
10 30.

131. An isolated nucleic acid molecule that exhibits at least 95% sequence
identity with a nucleic acid sequence selected from the group consisting of SEQ ID
NOS: 21 and 22, or at least 95% sequence identity with a fragment thereof comprising
15 at least 100 contiguous nucleotides thereof; and which further contains at least one
sequence that encodes a polypeptide which is at least 70% identical to a consensus
sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and
30.

20 132. An isolated nucleic acid molecule that exhibits about 95-99% sequence
identity with a nucleic acid sequence selected from the group consisting of SEQ ID
NOS: 21 and 22, or about 95-99% sequence identity with a fragment thereof
comprising at least 100 contiguous nucleotides thereof; and which further contains at
least one sequence that encodes a polypeptide which is at least 70% identical to a
25 consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27,
28, 29, and 30.

133. An isolated nucleic acid molecule having a nucleic acid sequence
selected from the group consisting of SEQ ID NOS: 21 and 22, or a fragment thereof
30 comprising at least 100 contiguous nucleotides thereof; and which further contains at
least one sequence that encodes a polypeptide which is at least 70% identical to a

consensus sequence selected from the group consisting of SEQ ID NOS: 25, 26, 27, 28, 29, and 30.

134. A method of screening a library comprising (a) hybridizing the library
5 with an isolated nucleic acid molecule according Claim 125 and (b) detecting one or more taste receptor clones in the library by a positive hybridization signal.

135. A method of detecting expression of a G Protein-Coupled Receptor polypeptide gene in a cell comprising:
10 contacting said cell with a nucleic acid molecule that hybridizes to an isolated nucleic acid molecule according to claim 125 under stringent conditions; and
detecting hybridization in order to detect expression of said G Protein-Coupled Receptor polypeptide gene.

136. A method for screening a mammalian genome for a coding sequence
15 for a G Protein-Coupled Receptor polypeptide active in taste signaling, comprising:

(i) contacting said mammalian genome with at least one degenerate primer consisting essentially of a nucleic acid sequence encoding a consensus sequence of SEQ ID NOS: 25, 26, 27, 28, 29, and 30, or at least one degenerate primer derived
20 from an isolated nucleic acid molecule according to claim 125;

(ii) amplifying said genomic sequence comprising said at least one primer sequence in the presence of polymerase, free nucleotides and cofactors; and

(iii) detecting the presence of an amplified sequence comprising a G Protein-Coupled Receptor polypeptide gene.

25 137. A biochemical assay for identifying tastant ligands having binding specificity for a G Protein-Coupled Receptor active in taste signaling, comprising:

(i) contacting one or more polypeptides according to claim 87 with a preparation of G Proteins and GTP γ S, and one or more putative tastant ligands or a
30 composition comprising one or more putative tastant ligands; and

(ii) detecting binding of a tastant ligand having binding specificity for said G Protein-Coupled Receptor active in taste signaling by measuring the binding of GTP γ S to the G Protein.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/10739

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/06719 A1 (THE REGENTS OF THE UNIVERSITY OF CALIFORNIA) 10 February 2000 (10/02/00), see entire document.	1-137

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

27 AUGUST 2001

Date of mailing of the international search report

20 SEP 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/10739

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

G01N 33/53, 33/567; C07K 14/47, 16/18, 14/705, 14/72; A61K 48/00; C12N 5/10, 15/12, 15/62, 15/63, 15/64

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

530/350, 387.1, 387.9, 388.22, 389.1; 536/23.1, 23.5, 24.3, 24.31; 435/6, 7.1, 7.2, 69.1, 69.7, 325, 471, 252.3, 254.11, 320.1; 800/21

B. FIELDS SEARCHED Minimum documentation searched Classification System: U.S.

530/350, 387.1, 387.9, 388.22, 389.1; 536/23.1, 23.5, 24.3, 24.31; 435/6, 7.1, 7.2, 69.1, 69.7, 325, 471, 252.3, 254.11, 320.1; 800/21

B. FIELDS SEARCHED Electronic data bases consulted (Name of data base and where practicable terms used):

search terms: taste-cell specific G Protein-Coupled Receptors, nucleic acid, polynucleotide, recombinant production, solid phase, screening, transgenic non-human, detection, assay, antibody

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/10739

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 1-137
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

all the claims recite specific SEQ ID NOS. However, since no Computer Readable Form (CRF) of the sequence listing has been filed in the case, and since the claims cannot be searched other than by a sequence search, the claims are unsearchable.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.